

# Symbolic control of stochastic systems via approximately bisimilar finite abstractions

Majid Zamani, Peyman Mohajerin Esfahani, Rupak Majumdar, Alessandro Abate, and John Lygeros

**Abstract**—Symbolic approaches to the control design over complex systems employ the construction of finite-state models that are related to the original control systems, then use techniques from finite-state synthesis to compute controllers satisfying specifications given in a temporal logic, and finally translate the synthesized schemes back as controllers for the concrete complex systems. Such approaches have been successfully developed and implemented for the synthesis of controllers over non-probabilistic control systems. In this paper, we extend the technique to probabilistic control systems modelled by controlled stochastic differential equations. We show that for every stochastic control system satisfying a probabilistic variant of incremental input-to-state stability, and for every given precision  $\varepsilon > 0$ , a finite-state transition system can be constructed, which is  $\varepsilon$ -approximately bisimilar to the original stochastic control system. Moreover, we provide results relating stochastic control systems to their corresponding finite-state transition systems in terms of probabilistic bisimulation relations known in the literature. We demonstrate the effectiveness of the construction by synthesizing controllers for stochastic control systems over rich specifications expressed in linear temporal logic. The discussed technique enables a new, automated, correct-by-construction controller synthesis approach for stochastic control systems, which are common mathematical models employed in many safety critical systems subject to structured uncertainty and are thus relevant for cyber-physical applications.

## I. INTRODUCTION, LITERATURE BACKGROUND, AND CONTRIBUTIONS

The design of controllers for complex control systems with respect to general temporal specifications in a reliable, yet cost-effective way, is a grand challenge in cyber-physical systems research. One promising direction is the use of *symbolic models*: symbolic models are discrete and finite approximations of the continuous dynamics constructed in a way that controllers designed for the approximations can be refined to controllers for the original dynamics. When finite symbolic models exist and can be constructed, one can leverage the apparatus of finite-state reactive synthesis [26] towards the problem of designing hybrid controllers. The formal notion of approximation is captured using  $\varepsilon$ -approximate bisimulation relations [15], which guarantee that

each trace of the continuous system can be matched by a trace of the symbolic model up to a precision  $\varepsilon$ , and vice versa.

The effective construction of finite symbolic models has been studied extensively for non-probabilistic control systems. Examples include work on piecewise-affine and multi-affine systems with constant control distribution [7], [17], abstractions based on convexity of reachable sets for sufficiently small sampling time [36], the use of incremental input-to-state stability [9], [16], [25], [34], [35], non-uniform abstractions of nonlinear systems over a finite-time horizon [40], and finally sound abstractions for unstable nonlinear control systems [45]. Together with automata-theoretic controller synthesis algorithms [26], effective symbolic models form the basis of controller synthesis tools such as Pessoa [27].

However, much less is known about continuous *stochastic* control systems. Existing results for probabilistic systems include the construction of finite abstractions for continuous-time stochastic dynamical systems under contractivity assumptions [1], for discrete-time stochastic hybrid dynamical systems endowed with certain continuity and ergodicity properties [3], and for discrete-time stochastic dynamical systems complying with a notion of bisimulation function [5]. While providing finite bisimilar abstractions, all the cited techniques are restricted to *autonomous* models (i.e., with no control inputs). As such, they are of interest for verification purposes, but fall short towards controller synthesis goals. On the other hand, for non-autonomous models there exist techniques to *check* if an infinite abstraction is formally related to a concrete stochastic control system via a notion of stochastic (bi)simulation function [19], however these results do not extend to the *construction* of approximations nor they deal with *finite* abstractions, and appear to be computationally intractable in the non-autonomous case because one requires to solve a game to compute stochastic (bi)simulation functions. Further, for specific temporal properties such as (probabilistic) invariance, there exist techniques to compute finite abstractions of discrete-time stochastic control systems [2], however their generalization to general properties in linear temporal logic is not obvious, nor their applicability to continuous-time models. The work in [38] provides algorithms for the verification and control problems restricted to probabilistic rectangular automata, in which random behaviors occur only over the discrete components – this limits their application to models with continuous probability laws. The work in [22] presents a finite Markov decision process approximation of a continuous-time linear stochastic control system for the verification of given temporal properties, however the relationship between abstract and concrete model is not quantitative. Along the same lines, classical discretization results in the literature [21] offer approximations of stochastic control systems that are related to the concrete models only asymptotically, rather than according

This work is supported by the European Commission MoVeS project FP7-ICT-2009-5 257005 and by a contract from Toyota.

M. Zamani is with the Department of Electrical Engineering and Information Technology, Technical University of Munich, D-80290 Munich, Germany. Email: zamani@tum.de.

P. Mohajerin Esfahani and J. Lygeros are with the Automatic Control Laboratory, ETH Zürich, ETL I22, 8092 Zürich, Switzerland. Email: {mohajerin, lygeros}@control.ee.ethz.ch.

R. Majumdar is with the Max Planck Institute for Software Systems, 67663 Kaiserslautern, Germany. Email: rupak@mpi-sws.org.

A. Abate is with the Department of Computer Science, University of Oxford, OX1 3QD Oxford, United Kingdom. Email: alessandro.abate@cs.ox.ac.uk.

to formal bisimulation or simulation notions that are in the end required to ensure the correspondence of controllers for temporal logic specifications over model trajectories.

In summary, to the best of our knowledge there is no comprehensive work on the construction of finite bisimilar abstractions for continuous-time continuous-space stochastic control systems. These systems offer a natural modeling framework for cyber-physical systems operating in an uncertain or noisy environment, and automated controller synthesis methodologies can enable more reliable system development at lower costs and times.

In this paper, we show the existence of  $\varepsilon$ -approximate bisimilar symbolic models (in the sense of moments) for continuous-time stochastic control systems satisfying a probabilistic version of the incremental input-to-state stability property [4], for any given parameter  $\varepsilon > 0$ . The symbolic models are finite if the continuous states lie within a bounded set. We also provide a simple way to construct the symbolic abstractions by quantizing the state and input sets. By guaranteeing the existence of an  $\varepsilon$ -approximate bisimulation relation among concrete and abstract models, we show that there exists a controller enforcing a desired specification on the symbolic model if and only if there exists a controller enforcing an  $\varepsilon$ -related specification on the original stochastic control system. The construction generalizes results for non-probabilistic systems [16], [25], [34], and reduces to these ones in the special case of dynamics with no noise. Moreover, it provides a framework upon which the new abstraction techniques can be developed for stochastic systems such as the recent results on stochastic control systems without any stability assumption [44]; on incrementally stable stochastic switched systems [42]; on incrementally stable randomly switched systems [43], and finally on incrementally stable stochastic control systems which do not require discretization of the set of states but only the set of inputs [47].

Furthermore, we provide quantitative results showing precisely how the proposed  $\varepsilon$ -approximate bisimulation relations among concrete stochastic control systems and abstract symbolic models are related to known probabilistic bisimulation notions recently developed in the literature. We also show the description of the proposed incremental stability in terms of a so-called incremental Lyapunov function. The derived notions are extensions of similar ones developed for non-probabilistic control systems [4], in the sense that the results for non-probabilistic control systems can be recovered in the framework of this paper by setting the diffusion term to zero.

We finally illustrate our results on two case studies, where controllers are synthesized over (non-linear) stochastic control systems with respect to rich linear temporal logic specifications of practical relevance.

## II. STOCHASTIC CONTROL SYSTEMS

### A. Notations

The identity map on a set  $A$  is denoted by  $1_A$ . If  $A$  is a subset of  $B$  we denote by  $\iota_A : A \hookrightarrow B$  or simply by  $\iota$  the natural inclusion map taking any  $a \in A$  to  $\iota(a) = a \in B$ . The symbols  $\mathbb{N}$ ,  $\mathbb{N}_0$ ,  $\mathbb{Z}$ ,  $\mathbb{R}$ ,  $\mathbb{R}^+$ , and  $\mathbb{R}_0^+$  denote the set of natural, nonnegative integer, integer, real, positive, and nonnegative real numbers, respectively. The symbols  $I_n$ ,  $0_n$ , and  $0_{n \times m}$  denote the identity matrix, the zero vector, and zero matrix in  $\mathbb{R}^{n \times n}$ ,  $\mathbb{R}^n$ , and  $\mathbb{R}^{n \times m}$ , respectively. Given a vector  $x \in \mathbb{R}^n$ ,

we denote by  $x_i$  the  $i$ -th element of  $x$ , and by  $\|x\|$  the infinity norm of  $x$ , namely  $\|x\| = \max\{|x_1|, |x_2|, \dots, |x_n|\}$ , where  $|x_i|$  denotes the absolute value of  $x_i$ . Given a matrix  $M = \{m_{ij}\} \in \mathbb{R}^{n \times m}$ , we denote by  $\|M\|$  the infinity norm of  $M$ , namely,  $\|M\| = \max_{1 \leq i \leq n} \sum_{j=1}^m |m_{ij}|$ , and by  $\|M\|_F$  the Frobenius norm of  $M$ , namely,  $\|M\|_F = \sqrt{\text{Tr}(MM^T)}$ , where  $\text{Tr}(P) = \sum_{i=1}^n p_{ii}$  for any  $P = \{p_{ij}\} \in \mathbb{R}^{n \times n}$ . We denote by  $\lambda_{\min}(A)$  and  $\lambda_{\max}(A)$  the minimum and maximum eigenvalues of matrix  $A$ , respectively. The diagonal set  $\Delta \subset \mathbb{R}^{2n}$  is defined as:  $\Delta = \{(x, x) \mid x \in \mathbb{R}^n\}$ . Given a set  $W \subseteq \mathbb{R}^n$  and  $\varepsilon \in \mathbb{R}_0^+$ , we denote by  $W^\varepsilon$  the  $\varepsilon$ -inflation of  $W$ , namely,  $W^\varepsilon = \{x \in \mathbb{R}^n \mid \|x\|_W \leq \varepsilon\}$ , where  $\|x\|_W = \inf_{w \in W} \|x - w\|$ .

The closed ball centered at  $x \in \mathbb{R}^n$  with radius  $\lambda$  is defined by  $\mathcal{B}_\lambda(x) = \{y \in \mathbb{R}^n \mid \|x - y\| \leq \lambda\}$ . A set  $B \subseteq \mathbb{R}^n$  is called a *box* if  $B = \prod_{i=1}^n [c_i, d_i]$ , where  $c_i, d_i \in \mathbb{R}$  with  $c_i < d_i$  for each  $i \in \{1, \dots, n\}$ . The *span* of a box  $B$  is defined as  $\text{span}(B) = \min\{|d_i - c_i| \mid i = 1, \dots, n\}$ . By defining  $[\mathbb{R}^n]_\eta = \{a \in \mathbb{R}^n \mid a_i = k_i \eta, k_i \in \mathbb{Z}, i = 1, \dots, n\}$ , the set  $\bigcup_{p \in [\mathbb{R}^n]_\eta} \mathcal{B}_\lambda(p)$  is a countable covering of  $\mathbb{R}^n$  for any  $\eta \in \mathbb{R}^+$  and  $\lambda \geq \eta/2$ . For a box  $B \subseteq \mathbb{R}^n$  and  $\eta \leq \text{span}(B)$ , define the  $\eta$ -approximation  $[B]_\eta = [\mathbb{R}^n]_\eta \cap B$ . Note that  $[B]_\eta \neq \emptyset$  for any  $\eta \leq \text{span}(B)$ . Geometrically, for any  $\eta \in \mathbb{R}^+$  with  $\eta \leq \text{span}(B)$  and  $\lambda \geq \eta$ , the collection of sets  $\{\mathcal{B}_\lambda(p)\}_{p \in [B]_\eta}$  is a finite covering of  $B$ , i.e.,  $B \subseteq \bigcup_{p \in [B]_\eta} \mathcal{B}_\lambda(p)$ . We extend the notions of *span* and of *approximation* to finite unions of boxes as follows. Let  $A = \bigcup_{j=1}^M A_j$ , where each  $A_j$  is a box. Define  $\text{span}(A) = \min\{\text{span}(A_j) \mid j = 1, \dots, M\}$ , and for any  $\eta \leq \text{span}(A)$ , define  $[A]_\eta = \bigcup_{j=1}^M [A_j]_\eta$ .

Given a measurable function  $f : \mathbb{R}_0^+ \rightarrow \mathbb{R}^n$ , the (essential) supremum of  $f$  is denoted by  $\|f\|_\infty$ ; we recall that  $\|f\|_\infty := (\text{ess})\sup\{\|f(t)\|, t \geq 0\}$ . A function  $f$  is essentially bounded if  $\|f\|_\infty < \infty$ . For a given time  $s \in \mathbb{R}^+$ , define  $f_s$  so that  $f_s(t) = f(t)$ , for any  $t \in [0, s]$ , and  $f_s(t) = 0$  elsewhere;  $f$  is said to be locally essentially bounded if for any  $s \in \mathbb{R}^+$ ,  $f_s$  is essentially bounded. A continuous function  $\gamma : \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$ , is said to belong to class  $\mathcal{K}$  if it is strictly increasing and  $\gamma(0) = 0$ ;  $\gamma$  is said to belong to class  $\mathcal{K}_\infty$  if  $\gamma \in \mathcal{K}$  and  $\gamma(r) \rightarrow \infty$  as  $r \rightarrow \infty$ . A continuous function  $\beta : \mathbb{R}_0^+ \times \mathbb{R}_0^+ \rightarrow \mathbb{R}_0^+$  is said to belong to class  $\mathcal{KL}$  if, for each fixed  $s$ , the map  $\beta(r, s)$  belongs to class  $\mathcal{K}$  with respect to  $r$  and, for each fixed nonzero  $r$ , the map  $\beta(r, s)$  is decreasing with respect to  $s$  and  $\beta(r, s) \rightarrow 0$  as  $s \rightarrow \infty$ . We identify a relation  $R \subseteq A \times B$  with the map  $R : A \rightarrow 2^B$  defined by  $b \in R(a)$  iff  $(a, b) \in R$ . Given a relation  $R \subseteq A \times B$ ,  $R^{-1}$  denotes the inverse relation defined by  $R^{-1} = \{(b, a) \in B \times A : (a, b) \in R\}$ . Given a twice continuously differentiable function  $V : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}_0^+$ , we denote by  $H(V)(x, x')$  the value of Hessian matrix of  $V$  at  $(x, x') \in \mathbb{R}^n \times \mathbb{R}^n$ .

### B. Stochastic control systems

Let  $(\Omega, \mathcal{F}, \mathbb{P})$  be a probability space endowed with a filtration  $\mathbb{F} = (\mathcal{F}_s)_{s \geq 0}$  satisfying the usual conditions of completeness and right continuity [30]. Let  $(W_s)_{s \geq 0}$  be a  $p$ -dimensional  $\mathbb{F}$ -Brownian motion.

*Definition 2.1:* A *stochastic control system* is a tuple  $\Sigma = (\mathbb{R}^n, \mathcal{U}, \mathcal{U}, f, \sigma)$ , where

- $\mathbb{R}^n$  is the state space;
- $\mathcal{U} \subseteq \mathbb{R}^m$  is an input set;

- $\mathcal{U}$  is a subset of the set of all measurable, locally essentially bounded functions of time from  $\mathbb{R}_0^+$  to  $\mathbb{U}$ ;
- $f : \mathbb{R}^n \times \mathbb{U} \rightarrow \mathbb{R}^n$  satisfies the following Lipschitz assumption: there exist constants  $L_x, L_u \in \mathbb{R}^+$  such that:  $\|f(x, u) - f(x', u')\| \leq L_x \|x - x'\| + L_u \|u - u'\|$  for all  $x, x' \in \mathbb{R}^n$  and all  $u, u' \in \mathbb{U}$ ;
- $\sigma : \mathbb{R}^n \rightarrow \mathbb{R}^{n \times p}$  satisfies the following Lipschitz assumption: there exists a constant  $Z \in \mathbb{R}_0^+$  such that:  $\|\sigma(x) - \sigma(x')\| \leq Z \|x - x'\|$  for all  $x, x' \in \mathbb{R}^n$ .

A stochastic process  $\xi : \Omega \times \mathbb{R}_0^+ \rightarrow \mathbb{R}^n$  is said to be a *solution process* of  $\Sigma$  if there exists  $v \in \mathcal{U}$  satisfying:

$$d\xi = f(\xi, v) dt + \sigma(\xi) dW_t, \quad (\text{II.1})$$

$\mathbb{P}$ -almost surely ( $\mathbb{P}$ -a.s.), where  $f$  is known as the drift,  $\sigma$  as the diffusion, and again  $W_t$  is Brownian motion. We also write  $\xi_{av}(t)$  to denote the value of the solution process at time  $t \in \mathbb{R}_0^+$  under the input signal  $v$  from initial condition  $\xi_{av}(0) = a$   $\mathbb{P}$ -a.s., in which  $a$  is a random variable that is measurable in  $\mathcal{F}_0$ . Let us remark that  $\mathcal{F}_0$  in general is not a trivial sigma-algebra, and thus the stochastic control system  $\Sigma$  can start from a random initial condition. Let us emphasize that the solution process is uniquely determined, since the assumptions on  $f$  and  $\sigma$  ensure its existence and uniqueness [30, Theorem 5.2.1, p. 68].

### III. A NOTION OF INCREMENTAL STABILITY

This section introduces a stability notion for stochastic control systems, which generalizes the notion of incremental input-to-state stability ( $\delta$ -ISS) [4] for non-probabilistic control systems. The main results presented in this work rely on the stability assumption discussed in this section.

#### A. Incremental stability and its description

*Definition 3.1:* A stochastic control system  $\Sigma$  is incrementally input-to-state stable in the  $q$ th moment ( $\delta$ -ISS- $M_q$ ), where  $q \geq 1$ , if there exist a  $\mathcal{KL}$  function  $\beta$  and a  $\mathcal{K}_\infty$  function  $\gamma$  such that for any  $t \in \mathbb{R}_0^+$ , any  $\mathbb{R}^n$ -valued random variables  $a$  and  $a'$  that are measurable in  $\mathcal{F}_0$ , and any  $v, v' \in \mathcal{U}$ , the following condition is satisfied:

$$\mathbb{E} [\|\xi_{av}(t) - \xi_{a'v'}(t)\|^q] \leq \beta(\mathbb{E} [\|a - a'\|^q], t) + \gamma(\|v - v'\|_\infty). \quad (\text{III.1})$$

It can be easily checked that a  $\delta$ -ISS- $M_q$  stochastic control system  $\Sigma$  is  $\delta$ -ISS in the absence of any noise as in the following:

$$\|\xi_{av}(t) - \xi_{a'v'}(t)\| \leq \beta(\|a - a'\|, t) + \gamma(\|v - v'\|_\infty), \quad (\text{III.2})$$

for  $a, a' \in \mathbb{R}^n$ , some  $\beta \in \mathcal{KL}$ , and some  $\gamma \in \mathcal{K}_\infty$ . Moreover, whenever  $f(0_n, 0_m) = 0_n$  and  $\sigma(0_n) = 0_{n \times p}$  (i.e., the drift and diffusion terms vanish at the origin), then  $\delta$ -ISS- $M_q$  implies input-to-state stability in the  $q$ th moment (ISS- $M_q$ ) [37] and global asymptotic stability in the  $q$ th moment (GAS- $M_q$ ) [10].

Similar to the use of  $\delta$ -ISS Lyapunov functions in the non-probabilistic case [4], we now describe  $\delta$ -ISS- $M_q$  in terms of the existence of *incremental Lyapunov functions*, as defined next.

*Definition 3.2:* Consider a stochastic control system  $\Sigma$  and a continuous function  $V : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}_0^+$  that is twice continuously differentiable on  $\{\mathbb{R}^n \times \mathbb{R}^n\} \setminus \Delta$ . The function  $V$  is called an incremental input-to-state stability in the  $q$ th

moment ( $\delta$ -ISS- $M_q$ ) Lyapunov function for  $\Sigma$ , where  $q \geq 1$ , if there exist  $\mathcal{K}_\infty$  functions  $\underline{\alpha}$ ,  $\bar{\alpha}$ ,  $\rho$ , and a constant  $\kappa \in \mathbb{R}^+$ , such that

- (i)  $\underline{\alpha}$  (resp.  $\bar{\alpha}$ ) is a convex (resp. concave) function;
- (ii) for any  $x, x' \in \mathbb{R}^n$ ,  $\underline{\alpha}(\|x - x'\|^q) \leq V(x, x') \leq \bar{\alpha}(\|x - x'\|^q)$ ;
- (iii) for any  $x, x' \in \mathbb{R}^n$ ,  $x \neq x'$ , and for any  $u, u' \in \mathbb{U}$ ,

$$\begin{aligned} \mathcal{L}^{u, u'} V(x, x') &:= [\partial_x V \quad \partial_{x'} V] \begin{bmatrix} f(x, u) \\ f(x', u') \end{bmatrix} \\ &+ \frac{1}{2} \text{Tr} \left( \begin{bmatrix} \sigma(x) \\ \sigma(x') \end{bmatrix} \begin{bmatrix} \sigma^T(x) & \sigma^T(x') \end{bmatrix} \begin{bmatrix} \partial_{x, x} V & \partial_{x, x'} V \\ \partial_{x', x} V & \partial_{x', x'} V \end{bmatrix} \right) \\ &\leq -\kappa V(x, x') + \rho(\|u - u'\|), \end{aligned}$$

where  $\mathcal{L}^{u, u'}$  is the infinitesimal generator associated to the stochastic control system (II.1) [30, Section 7.3], which in this case depends on two separate controls  $u, u'$ . The symbols  $\partial_x$  and  $\partial_{x, x'}$  denote first- and second-order partial derivatives with respect to  $x$  and  $x'$ , respectively.

Roughly speaking, condition (ii) implies that the growth rate of functions  $\bar{\alpha}$  and  $\underline{\alpha}$  are linear, as a concave function is supposed to dominate a convex one. However, these conditions do not restrict the behavior of  $\bar{\alpha}$  and  $\underline{\alpha}$  to only linear functions on a compact subset of  $\mathbb{R}^n$ . Note that condition (i) is not required in the context of non-probabilistic control systems. The following theorem clarifies why such a requirement is instead necessary for a stochastic control system, and describes  $\delta$ -ISS- $M_q$  in terms of the existence of  $\delta$ -ISS- $M_q$  Lyapunov functions.

*Theorem 3.3:* A stochastic control system  $\Sigma$  is  $\delta$ -ISS- $M_q$  if it admits a  $\delta$ -ISS- $M_q$  Lyapunov function.

The proof of Theorem 3.3 is provided in the Appendix.

One can resort to available software tools, such as SOS-TOOLS [31, Subsection 4.2], to search for appropriate, non-trivial  $\delta$ -ISS- $M_q$  Lyapunov functions of the form of polynomial. Satisfaction of conditions (i) and (ii) of Definition 3.2 globally on  $\mathbb{R}^n$  may require  $\underline{\alpha}$  and  $\bar{\alpha}$  to be piecewise polynomial functions. However, as long as one is interested in the dynamics of  $\Sigma$  on a compact subset of  $\mathbb{R}^n$ , those conditions can be still satisfied by  $\underline{\alpha}$  and  $\bar{\alpha}$  of the form of polynomial.

Now we look into special instances where function  $V$  can be easily computed based on the model dynamics. The first result provides a sufficient condition for a particular function  $V$  to be a  $\delta$ -ISS- $M_q$  Lyapunov function for a stochastic control system  $\Sigma$ , when  $q = 1, 2$  (that is, in the first or second moment).

*Lemma 3.4:* Consider a stochastic control system  $\Sigma$ . Let  $q \in \{1, 2\}$ ,  $P \in \mathbb{R}^{n \times n}$  be a symmetric positive definite matrix, and the function  $V : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}_0^+$  be defined as follows:

$$V(x, x') := \left( \tilde{V}(x, x') \right)^{\frac{2}{q}} = \left( \frac{1}{q} (x - x')^T P (x - x') \right)^{\frac{2}{q}}, \quad (\text{III.3})$$

and satisfy

$$\begin{aligned} (x - x')^T P (f(x, u) - f(x', u)) + \frac{1}{2} \left\| \sqrt{P} (\sigma(x) - \sigma(x')) \right\|_F^2 \\ \leq -\tilde{\kappa} (V(x, x'))^{\frac{2}{q}}, \end{aligned} \quad (\text{III.4})$$

or, if  $f$  is differentiable with respect to  $x$ , satisfy

$$\begin{aligned} (x - x')^T P \partial_x f(z, u) (x - x') + \frac{1}{2} \left\| \sqrt{P} (\sigma(x) - \sigma(x')) \right\|_F^2 \\ \leq -\tilde{\kappa} (V(x, x'))^{\frac{2}{q}}, \end{aligned} \quad (\text{III.5})$$

for all  $x, x', z \in \mathbb{R}^n$ , for all  $u \in \mathcal{U}$ , and for some constant  $\tilde{\kappa} \in \mathbb{R}^+$ . Then  $V$  is a  $\delta$ -ISS- $M_q$  Lyapunov function for  $\Sigma$ .

The proof of Lemma 3.4 is provided in the Appendix.

The next result provides a condition that is equivalent to (III.4) or (III.5) for linear stochastic control systems  $\Sigma$  (that is, for systems with linear drift and diffusion terms) in the form of a linear matrix inequality (LMI), which can be easily solved numerically.

*Corollary 3.5:* Consider a stochastic control system  $\Sigma$ , where for all  $x \in \mathbb{R}^n$ , and  $u \in \mathcal{U}$ ,  $f(x, u) := Ax + Bu$ , for some  $A \in \mathbb{R}^{n \times n}$ ,  $B \in \mathbb{R}^{n \times m}$ , and  $\sigma(x) := [\sigma_1 x \ \sigma_2 x \ \cdots \ \sigma_p x]$ , for some  $\sigma_i \in \mathbb{R}^{n \times n}$ . Then, function  $V$  in (III.3) is a  $\delta$ -ISS- $M_q$  Lyapunov function for  $\Sigma$ , when  $q \in \{1, 2\}$ , if there exists a constant  $\hat{\kappa} \in \mathbb{R}^+$  satisfying the following LMI:

$$PA + A^T P + \sum_{i=1}^p \sigma_i^T P \sigma_i \preceq -\hat{\kappa} P. \quad (\text{III.6})$$

*Proof:* The corollary is a particular case of Lemma 3.4. It suffices to show that for linear dynamics, the LMI in (III.6) yields the condition in (III.4) or (III.5). First it is straightforward to observe that

$$\begin{aligned} \|\sqrt{P}(\sigma(x) - \sigma(x'))\|_F^2 &= \text{Tr}((\sigma(x) - \sigma(x'))^T P (\sigma(x) - \sigma(x'))) \\ &= (x - x')^T \sum_{i=1}^p \sigma_i^T P \sigma_i (x - x'), \end{aligned}$$

and that

$$(x - x')^T P \partial_x f(z, u) (x - x') = \frac{1}{2} (x - x')^T (PA + A^T P) (x - x'),$$

for any  $x, x', z \in \mathbb{R}^n$  and any  $u \in \mathcal{U}$ . Now suppose there exists  $\hat{\kappa} \in \mathbb{R}^+$  such that (III.6) holds. It can be readily verified that the desired requirements in (III.4) and (III.5) are verified by choosing  $\tilde{\kappa} = \frac{q\hat{\kappa}}{2}$ . ■

*Remark 3.6:* By considering both  $\hat{\kappa}$  and  $P$  as decision variables, inequality (III.6) is not an LMI anymore as the right hand side is bilinear. However, one may resort to iterative schemes (e.g. bisection algorithm on scalar variable  $\hat{\kappa}$ ) to optimize over the parameters  $\hat{\kappa}$  and  $P$ , separately. Note that seeking appropriate matrix  $P$  and constant  $\hat{\kappa}$  can provide tighter upper bounds in (III.1).

*Remark 3.7:* Consider a stochastic control system  $\Sigma$ . Assume that  $f$  is differentiable with respect to  $x$  and that, for any  $x \in \mathbb{R}^n$ ,  $\sigma(x) := [\sigma_1 x \ \sigma_2 x \ \cdots \ \sigma_p x]$ , for some  $\sigma_i \in \mathbb{R}^{n \times n}$ . Then, the function  $V$  in (III.3) is a  $\delta$ -ISS- $M_q$  Lyapunov function for  $\Sigma$ , when  $q \in \{1, 2\}$ , if there exists a constant  $\hat{\kappa} \in \mathbb{R}^+$  satisfying the following matrix inequality:

$$P \partial_x f(z, u) + (\partial_x f(z, u))^T P + \sum_{i=1}^p \sigma_i^T P \sigma_i \preceq -\hat{\kappa} P, \quad (\text{III.7})$$

for any  $z \in \mathbb{R}^n$  and any  $u \in \mathcal{U}$ . One can easily verify that condition (III.7) corresponds to the contractivity conditions (with respect to the states) in [46], obtained with contraction metric  $P$ , and to the Demidovich's condition in [32] for a system  $\Sigma$  in the absence of any noise, i.e.  $\sigma_i = 0_{n \times n}$  for all  $i = 1, \dots, p$ . □

## B. Noisy and noise-free trajectories

In order to introduce a symbolic model in Section V for the stochastic control system  $\Sigma$ , we need the following technical results, which provide an upper bound on the distance (in

the  $q$ th moment) between the solution processes of  $\Sigma$  and those of the corresponding non-probabilistic control system, i.e.  $\bar{\Sigma} = (\mathbb{R}^n, \mathcal{U}, \mathcal{U}, f, 0_{n \times p})$ , obtained by disregarding the diffusion term (that is,  $\sigma$ ). From now on, we use the notation  $\xi_{xv}$  to denote the trajectory of  $\bar{\Sigma}$  under the input signal  $v$  from initial condition  $x$ .

*Lemma 3.8:* Consider a stochastic control system  $\Sigma$  such that  $f(0_n, 0_m) = 0_n$  and  $\sigma(0_n) = 0_{n \times p}$ . Suppose that  $q \geq 2$  and there exists a  $\delta$ -ISS- $M_q$  Lyapunov function  $V$  for  $\Sigma$  such that its Hessian is a positive semidefinite matrix in  $\mathbb{R}^{2n \times 2n}$  and  $\partial_{x,x} V(x, x') \leq P$ , for any  $x, x' \in \mathbb{R}^n$  and some positive semidefinite matrix  $P \in \mathbb{R}^{n \times n}$ . Then for any  $x$  in a compact set  $D \subset \mathbb{R}^n$  and any  $v \in \mathcal{U}$ , we have

$$\mathbb{E} \left[ \|\xi_{xv}(t) - \bar{\xi}_{xv}(t)\|^q \right] \leq h(\sigma, t), \quad (\text{III.8})$$

where the nonnegative valued function  $h$  tends to zero as  $t \rightarrow 0$ ,  $t \rightarrow +\infty$ , or as  $Z \rightarrow 0$ , where  $Z$  is the Lipschitz constant, introduced in Definition 2.1.

The proof of Lemma 3.8 is provided in the Appendix.

One can compute explicitly the function  $h$  using equation (IX.7) in the Appendix. Although function  $h$  in (IX.7) is only an explicit function of time  $t$ , with a slight abuse of notation, we prefer to keep  $\sigma$  as one of the arguments of  $h$  to show later how the results in non-probabilistic cases in the literature can be recovered by setting  $\sigma = 0$  in function  $h$ .

*Remark 3.9:* In the previous lemma, if  $V$  is a polynomial, then the condition  $H(V)(x, x') \succeq 0_{2n \times 2n}$  for all  $x, x' \in \mathbb{R}^n$  is equivalent to  $V$  being a convex function [11]. Furthermore, if we assume that for all  $x, x' \in \mathbb{R}^n$ ,  $H(V)(x, x') = M(x, x')^T M(x, x')$ , where  $M(x, x') \in \mathbb{R}^{s \times 2n}$  is a polynomial matrix for some  $s \in \mathbb{N}$ , then  $V$  is a sum of squares which can be efficiently searched through convex linear matrix inequalities optimizations [11], and using some available tools, such as SOSTOOLS [31, Subsection 4.2]. □

The following lemma provides an explicit result in line with that of Lemma 3.8 for a model  $\Sigma$  admitting a  $\delta$ -ISS- $M_q$  Lyapunov function  $V$  as in (III.3), where  $q \in \{1, 2\}$ .

*Lemma 3.10:* Consider a stochastic control system  $\Sigma$  such that  $f(0_n, 0_m) = 0_n$  and  $\sigma(0_n) = 0_{n \times p}$ . Suppose that the function  $V$  in (III.3) satisfies (III.4) or (III.5) for  $\Sigma$ . For any  $x$  in a compact set  $D \subset \mathbb{R}^n$  and any  $v \in \mathcal{U}$ , we have  $\mathbb{E} \left[ \|\xi_{xv}(t) - \bar{\xi}_{xv}(t)\|^2 \right] \leq h(\sigma, t)$ , where

$$\begin{aligned} h(\sigma, t) &= \frac{2 \|\sqrt{P}\|^2 n^2 \min\{n, p\} Z^2 e^{-\frac{2\tilde{\kappa}t}{q}}}{\lambda_{\min}^2(P) \tilde{\kappa}} \\ &\left( \lambda_{\max}(P) \left(1 - e^{-\frac{\tilde{\kappa}t}{2}}\right) \sup_{x \in D} \{\|x\|^2\} + \frac{\|\sqrt{P}\|^2 L_u^2}{e\tilde{\kappa}} \sup_{u \in \mathcal{U}} \{\|u\|^2\} t \right), \end{aligned}$$

$Z$  is the Lipschitz constant, as introduced in Definition 2.1. It can be readily verified that the nonnegative valued function  $h$  tends to zero as  $t \rightarrow 0$ ,  $t \rightarrow +\infty$ , or as  $Z \rightarrow 0$ .

The proof of Lemma 3.10 is provided in the Appendix.

For a linear stochastic control system  $\Sigma$ , the following corollary tailors the result in Lemma 3.10 obtaining possibly a less conservative expression for function  $h$ , based on parameters of drift and diffusion.

*Corollary 3.11:* Consider a stochastic control system  $\Sigma$ , where for all  $x \in \mathbb{R}^n$  and  $u \in \mathbb{R}^m$ ,  $f(x) := Ax + Bu$ , for some  $A \in \mathbb{R}^{n \times n}$ ,  $B \in \mathbb{R}^{n \times m}$ , and  $\sigma(x) := [\sigma_1 x \ \sigma_2 x \ \cdots \ \sigma_p x]$ , for some  $\sigma_i \in \mathbb{R}^{n \times n}$ . Suppose that the function  $V$  in (III.3) satisfies (III.6) for  $\Sigma$ . For any  $x$  in a compact set  $D$  and any

$v \in \mathcal{U}$ , we have  $\mathbb{E} \left[ \left\| \xi_{xv}(t) - \bar{\xi}_{xv}(t) \right\|^2 \right] \leq h(\sigma, t)$ , where

$$h(\sigma, t) = \frac{n\lambda_{\max} \left( \sum_{i=1}^p \sigma_i^T P \sigma_i \right) e^{-\hat{\kappa}t}}{\lambda_{\min}(P)} \int_0^t \left( \left\| e^{As} \right\| \sup_{x \in \mathbb{D}} \{ \|x\| \} + \left( \int_0^s \left\| e^{Ar} B \right\| dr \right) \sup_{u \in \mathcal{U}} \{ \|u\| \} \right)^2 ds.$$

The proof of Corollary 3.11 is provided in the Appendix.

#### IV. SYMBOLIC MODELS

##### A. Systems

We employ the notion of system [39] to describe both stochastic control systems as well as their symbolic models.

*Definition 4.1:* A system  $S$  is a tuple  $S = (X, X_0, U, \longrightarrow, Y, H)$ , where  $X$  is a set of states,  $X_0 \subseteq X$  is a set of initial states,  $U$  is a set of inputs,  $\longrightarrow \subseteq X \times U \times X$  is a transition relation,  $Y$  is a set of outputs, and  $H : X \rightarrow Y$  is an output map.

A transition  $(x, u, x') \in \longrightarrow$  is also denoted by  $x \xrightarrow{u} x'$ . For a transition  $x \xrightarrow{u} x'$ , state  $x'$  is called a  $u$ -successor, or simply a successor, of state  $x$ . For technical reasons, we assume that for any  $x \in X$ , there exists some  $u$ -successor of  $x$ , for some  $u \in U$  – let us remark that this is always the case for the considered systems later in this paper.

System  $S$  is said to be

- *metric*, if the output set  $Y$  is equipped with a metric  $\mathbf{d} : Y \times Y \rightarrow \mathbb{R}_0^+$ ;
- *finite* (or *symbolic*), if  $X$  and  $U$  are finite sets;
- *deterministic*, if for any state  $x \in X$  and any input  $u$ , there exists at most one  $u$ -successor.

For a system  $S = (X, X_0, U, \longrightarrow, Y, H)$  and given any initial state  $x_0 \in X_0$ , a finite state run generated from  $x_0$  is a finite sequence of transitions:  $x_0 \xrightarrow{u_0} x_1 \xrightarrow{u_1} x_2 \xrightarrow{u_2} \dots \xrightarrow{u_{n-2}} x_{n-1} \xrightarrow{u_{n-1}} x_n$ , such that  $x_i \xrightarrow{u_i} x_{i+1}$  for all  $0 \leq i < n$ . A finite state run can be directly extended to an infinite state run as well.

##### B. System relations

We recall the notion of approximate (bi)simulation relation, introduced in [15], which is useful when analyzing or synthesizing controllers for deterministic systems.

*Definition 4.2:* Let  $S_a = (X_a, X_{a0}, U_a, \xrightarrow{a}, Y_a, H_a)$  and  $S_b = (X_b, X_{b0}, U_b, \xrightarrow{b}, Y_b, H_b)$  be metric systems with the same output sets  $Y_a = Y_b$  and metric  $\mathbf{d}$ . For  $\varepsilon \in \mathbb{R}_0^+$ , a relation  $R \subseteq X_a \times X_b$  is said to be an  $\varepsilon$ -approximate simulation relation from  $S_a$  to  $S_b$  if the following three conditions are satisfied:

- (i) for every  $x_{a0} \in X_{a0}$ , there exists  $x_{b0} \in X_{b0}$  with  $(x_{a0}, x_{b0}) \in R$ ;
- (ii) for every  $(x_a, x_b) \in R$ , we have  $\mathbf{d}(H_a(x_a), H_b(x_b)) \leq \varepsilon$ ;
- (iii) for every  $(x_a, x_b) \in R$ , we have that  $x_a \xrightarrow{u_a} x'_a$  in  $S_a$

implies the existence of  $x_b \xrightarrow{u_b} x'_b$  in  $S_b$  satisfying  $(x'_a, x'_b) \in R$ .

A relation  $R \subseteq X_a \times X_b$  is said to be an  $\varepsilon$ -approximate bisimulation relation between  $S_a$  and  $S_b$  if  $R$  is an  $\varepsilon$ -approximate

simulation relation from  $S_a$  to  $S_b$  and  $R^{-1}$  is an  $\varepsilon$ -approximate simulation relation from  $S_b$  to  $S_a$ .

System  $S_a$  is  $\varepsilon$ -approximately simulated by  $S_b$ , or  $S_b$   $\varepsilon$ -approximately simulates  $S_a$ , denoted by  $S_a \preceq_{\mathcal{S}}^{\varepsilon} S_b$ , if there exists an  $\varepsilon$ -approximate simulation relation from  $S_a$  to  $S_b$ . System  $S_a$  is  $\varepsilon$ -approximate bisimilar to  $S_b$ , denoted by  $S_a \cong_{\mathcal{S}}^{\varepsilon} S_b$ , if there exists an  $\varepsilon$ -approximate bisimulation relation between  $S_a$  and  $S_b$ .

Note that when  $\varepsilon = 0$ , condition (ii) in the above definition changes as  $(x_a, x_b) \in R$  if and only if  $H_a(x_a) = H_b(x_b)$ , and  $R$  becomes an exact simulation relation, as introduced in [29]. Similarly, whenever  $\varepsilon = 0$ ,  $R$  becomes an exact bisimulation relation.

#### V. SYMBOLIC MODELS FOR STOCHASTIC CONTROL SYSTEMS

This section contains the main contribution of the paper. We show that for any stochastic control system  $\Sigma$  admitting a  $\delta$ -ISS- $M_q$  Lyapunov function, and for any precision level  $\varepsilon \in \mathbb{R}^+$ , we can construct a finite system that is  $\varepsilon$ -approximately bisimilar to  $\Sigma$ . In order to do so, we use the notion of system as an abstract representation of a stochastic control system, capturing all the information contained in it. More precisely, given a stochastic control system  $\Sigma$ , we define an associated metric system  $S(\Sigma) = (X, X_0, U, \longrightarrow, Y, H)$ , where:

- $X$  is the set of all  $\mathbb{R}^n$ -valued random variables defined on the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ ;
- $X_0$  is the set of all  $\mathbb{R}^n$ -valued random variables that are measurable over the trivial sigma-algebra  $\mathcal{F}_0$ , i.e., the system starts from a non-probabilistic initial condition, which is equivalently a random variable with a Dirac probability distribution;
- $U = \mathcal{U}$ ;
- $x \xrightarrow{v} x'$  if  $x$  and  $x'$  are measurable in  $\mathcal{F}_t$  and  $\mathcal{F}_{t+\tau}$ , respectively, for some  $t \in \mathbb{R}_0^+$  and  $\tau \in \mathbb{R}^+$ , and there exists a solution process  $\xi : \Omega \times \mathbb{R}_0^+ \rightarrow \mathbb{R}^n$  of  $\Sigma$  satisfying  $\xi(t) = x$  and  $\xi_{xv}(\tau) = x'$  P-a.s.;
- $Y$  is the set of all  $\mathbb{R}^n$ -valued random variables defined on the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ ;
- $H = 1_X$ .

We assume that the output set  $Y$  is equipped with the metric  $\mathbf{d}(y, y') = (\mathbb{E} [\|y - y'\|^q])^{\frac{1}{q}}$ , for any  $y, y' \in Y$  and some  $q \geq 1$ . Let us remark that the set of states of  $S(\Sigma)$  is uncountable and that  $S(\Sigma)$  is a deterministic system in the sense of Definition 4.1, since (cf. Subsection II-B) the solution process of  $\Sigma$  is uniquely determined.

The results in this section rely on additional assumptions on model  $\Sigma$  that are described next (however such assumptions are not required for the definitions and results in Sections II, III, and IV). We restrict our attention to stochastic control systems  $\Sigma$  with  $f(0_n, 0_m) = 0_n$ ,  $\sigma(0_n) = 0_{n \times p}$ , and input sets  $\mathcal{U}$  that are assumed to be finite unions of boxes. We further restrict our attention to sampled-data stochastic control systems, where input signals belong to set  $\mathcal{U}_{\tau}$  which contains only signals that are constant over intervals of length  $\tau \in \mathbb{R}^+$ , i.e.

$$\mathcal{U}_{\tau} = \left\{ v \in \mathcal{U} \mid v(t) = v((k-1)\tau), t \in [(k-1)\tau, k\tau], k \in \mathbb{N} \right\}.$$

Let us denote by  $S_{\tau}(\Sigma)$  a sub-system of  $S(\Sigma)$  obtained by selecting those transitions of  $S(\Sigma)$  corresponding to solution processes of duration  $\tau$  and to control inputs in  $\mathcal{U}_{\tau}$ . This can be seen as the time discretization or as the sampling of

a process. More precisely, given a stochastic control system  $\Sigma = (\mathbb{R}^n, \mathbf{U}, \mathcal{U}_\tau, f, \sigma)$ , we define the associated metric system  $S_\tau(\Sigma) = \left( X_\tau, X_{\tau_0}, U_\tau, \xrightarrow{\tau}, Y_\tau, H_\tau \right)$ , where  $X_\tau = X$ ,  $X_{\tau_0} = X_0$ ,  $U_\tau = \mathcal{U}_\tau$ ,  $Y_\tau = \hat{Y}$ ,  $H_\tau = H$ , and

- $x_\tau \xrightarrow{v_\tau} x'_\tau$  if  $x_\tau$  and  $x'_\tau$  are measurable, respectively, in  $\mathcal{F}_{k\tau}$  and  $\mathcal{F}_{(k+1)\tau}$  for some  $k \in \mathbb{N}_0$ , and there exists a solution process  $\xi : \Omega \times \mathbb{R}_0^+ \rightarrow \mathbb{R}^n$  of  $\Sigma$  satisfying  $\xi(k\tau) = x_\tau$  and  $\xi_{x_\tau v_\tau}(\tau) = x'_\tau$  P-a.s..

Notice that a finite state run  $x_0 \xrightarrow{v_0} x_1 \xrightarrow{v_1} \dots \xrightarrow{v_{N-1}} x_N$ , of  $S_\tau(\Sigma)$ , where  $v_i \in \mathcal{U}_\tau$  and  $x_i = \xi_{x_{i-1} v_{i-1}}(\tau)$  P-a.s. for  $i = 1, \dots, N$ , captures the solution process of the stochastic control system  $\Sigma$  at times  $t = 0, \tau, \dots, N\tau$ , started from the non-probabilistic initial condition  $x_0$  and resulting from a control input  $v$  obtained by the concatenation of the input signals  $v_i$  (i.e.  $v(t) = v_{i-1}(t)$  for any  $t \in [(i-1)\tau, i\tau[$ ), for  $i = 1, \dots, N$ .

Let us proceed introducing a fully symbolic system for the concrete model  $\Sigma$ . Consider a stochastic control system  $\Sigma = (\mathbb{R}^n, \mathbf{U}, \mathcal{U}_\tau, f, \sigma)$  and a triple  $\mathbf{q} = (\tau, \eta, \mu)$  of quantization parameters, where  $\tau$  is the sampling time,  $\eta$  is the state space quantization, and  $\mu$  is the input set quantization. Given  $\Sigma$  and  $\mathbf{q}$ , consider the following system:

$$S_{\mathbf{q}}(\Sigma) = (X_{\mathbf{q}}, X_{\mathbf{q}0}, U_{\mathbf{q}}, \xrightarrow{\mathbf{q}}, Y_{\mathbf{q}}, H_{\mathbf{q}}), \quad (\text{V.1})$$

consisting of (cf. Notation in Subsection II-A):

- $X_{\mathbf{q}} = \llbracket \mathbb{R}^n \rrbracket_\eta$ ;
- $X_{\mathbf{q}0} = \llbracket \mathbb{R}^n \rrbracket_\eta$ ;
- $U_{\mathbf{q}} = \llbracket \mathbf{U} \rrbracket_\mu$ ;
- $x_{\mathbf{q}} \xrightarrow{u_{\mathbf{q}}} x'_{\mathbf{q}}$  if there exists a  $x'_q \in X_{\mathbf{q}}$  such that  $\left\| \bar{\xi}_{x_{\mathbf{q}} u_{\mathbf{q}}}(\tau) - x'_q \right\| \leq \eta$ ;
- $Y_{\mathbf{q}}$  is the set of all  $\mathbb{R}^n$ -valued random variables defined on the probability space  $(\Omega, \mathcal{F}, \mathbb{P})$ ;
- $H_{\mathbf{q}} = \iota : X_{\mathbf{q}} \hookrightarrow Y_{\mathbf{q}}$ .

Note that we have abused notation by identifying  $u_{\mathbf{q}} \in \llbracket \mathbf{U} \rrbracket_\mu$  with the constant input signal with domain  $[0, \tau[$  and value  $u_{\mathbf{q}}$ . Notice that the proposed abstraction  $S_{\mathbf{q}}(\Sigma)$  is a deterministic system in the sense of Definition 4.1. In order to establish an approximate bisimulation relation, the output set  $Y_{\mathbf{q}}$  is defined similarly to the stochastic system  $S_\tau(\Sigma)$ . Therefore, in the definition of  $H_{\mathbf{q}}$ , the inclusion map  $\iota$  is meant, with a slight abuse of notation, as a mapping from a non-probabilistic grid point to a random variable with a Dirac probability distribution centered at that grid point. As argued in [39], there is no loss of generality to alternatively assume that  $Y_{\mathbf{q}} = X_{\mathbf{q}}$  and  $H_{\mathbf{q}} = 1_{X_{\mathbf{q}}}$ . For later use, we denote by  $X_{x_{\mathbf{q}0} u} : \mathbb{N}_0 \rightarrow X_{\mathbf{q}}$  a state run of  $S_{\mathbf{q}}(\Sigma)$  from initial condition  $X_{x_{\mathbf{q}0} u}(0) = x_{\mathbf{q}0}$  and under input sequence  $u : \mathbb{N}_0 \rightarrow \llbracket \mathbf{U} \rrbracket_\mu$ .

The transition relation of  $S_{\mathbf{q}}(\Sigma)$  is well defined in the sense that for every  $x_{\mathbf{q}} \in \llbracket \mathbb{R}^n \rrbracket_\eta$  and every  $u_{\mathbf{q}} \in \llbracket \mathbf{U} \rrbracket_\mu$  there always exists  $x'_q \in \llbracket \mathbb{R}^n \rrbracket_\eta$  such that  $x_{\mathbf{q}} \xrightarrow{u_{\mathbf{q}}} x'_q$ . This can be seen since by definition of  $\llbracket \mathbb{R}^n \rrbracket_\eta$ , for any  $\hat{x} \in \mathbb{R}^n$  there always exists a state  $\hat{x}' \in \llbracket \mathbb{R}^n \rrbracket_\eta$  such that  $\|\hat{x} - \hat{x}'\| \leq \eta$ . Hence, for  $\bar{\xi}_{x_{\mathbf{q}} u_{\mathbf{q}}}(\tau)$  there always exists a state  $x'_q \in \llbracket \mathbb{R}^n \rrbracket_\eta$  satisfying  $\left\| \bar{\xi}_{x_{\mathbf{q}} u_{\mathbf{q}}}(\tau) - x'_q \right\| \leq \eta$ .

## A. Main results

In order to show the first main result of this work, we raise a supplementary assumption on the  $\delta$ -ISS- $M_{\mathbf{q}}$  Lyapunov function  $V$  as follows:

$$|V(x, y) - V(x, z)| \leq \hat{\gamma}(\|y - z\|), \quad (\text{V.2})$$

for any  $x, y, z \in \mathbb{R}^n$ , and some  $\mathcal{K}_\infty$  and concave function  $\hat{\gamma}$ . This assumption is not restrictive, provided  $V$  is restricted to a compact subset of  $\mathbb{R}^n \times \mathbb{R}^n$ . Indeed, for all  $x, y, z \in D$ , where  $D \subset \mathbb{R}^n$  is compact, by applying the mean value theorem to the function  $y \rightarrow V(x, y)$ , one gets

$$|V(x, y) - V(x, z)| \leq \hat{\gamma}(\|y - z\|),$$

$$\text{where } \hat{\gamma}(r) = \left( \max_{x, y \in D \setminus \Delta} \left\| \frac{\partial V(x, y)}{\partial y} \right\| \right) r.$$

In particular, for the  $\delta$ -ISS- $M_1$  Lyapunov function  $V$  defined in (III.3), we obtain explicitly that  $\hat{\gamma}(r) = \frac{\lambda_{\max}(P)}{\sqrt{\lambda_{\min}(P)}} r$  [39, Proposition 10.5] satisfying (V.2) globally over  $\mathbb{R}^n \times \mathbb{R}^n$ . We can now present the first main result of the paper, which relates the existence of a  $\delta$ -ISS- $M_{\mathbf{q}}$  Lyapunov function to the construction of a symbolic model.

*Theorem 5.1:* Consider a stochastic control system  $\Sigma$ , admitting a  $\delta$ -ISS- $M_{\mathbf{q}}$  Lyapunov function  $V$ , of the form of (III.3) or the one explained in Lemma 3.8. For any  $\varepsilon \in \mathbb{R}^+$  and any triple  $\mathbf{q} = (\tau, \eta, \mu)$  of quantization parameters satisfying  $\mu \leq \text{span}(\mathbf{U})$  and

$$\bar{\alpha}(\eta^q) \leq \underline{\alpha}(\varepsilon^q), \quad (\text{V.3})$$

$$e^{-\kappa\tau} \underline{\alpha}(\varepsilon^q) + \frac{1}{e\kappa} \rho(\mu) + \hat{\gamma} \left( (h(\sigma, \tau))^{\frac{1}{q}} + \eta \right) \leq \underline{\alpha}(\varepsilon^q), \quad (\text{V.4})$$

we have that  $S_{\mathbf{q}}(\Sigma) \cong_{\mathcal{S}}^{\varepsilon} S_\tau(\Sigma)$ .

The proof of Theorem 5.1 is provided in the Appendix.

It can be readily seen that when we are interested in the dynamics of  $\Sigma$ , initialized on a compact  $D \subset \mathbb{R}^n$  of the form of finite union of boxes and for a given precision  $\varepsilon$ , there always exists a sufficiently large value of  $\tau$  and small enough values of  $\eta$  and  $\mu$ , such that  $\eta \leq \text{span}(D)$  and such that the conditions in (V.3) and (V.4) are satisfied. On the other hand, for a given fixed sampling time  $\tau$ , one can find sufficiently small values of  $\eta$  and  $\mu$  satisfying  $\eta \leq \text{span}(D)$ , (V.3) and (V.4), as long as the precision  $\varepsilon$  is lower bounded by:

$$\varepsilon > \left( \underline{\alpha}^{-1} \left( \frac{\hat{\gamma} \left( (h(\sigma, \tau))^{\frac{1}{q}} \right)}{1 - e^{-\kappa\tau}} \right) \right)^{\frac{1}{q}}. \quad (\text{V.5})$$

One can easily verify that the lower bound on  $\varepsilon$  in (V.5) goes to zero as  $\tau$  goes to infinity or as  $Z \rightarrow 0$ , where  $Z$  is the Lipschitz constant, introduced in Definition 2.1. Furthermore, one can try to minimize the lower bound on  $\varepsilon$  in (V.5) by appropriately choosing a  $\delta$ -ISS- $M_{\mathbf{q}}$  Lyapunov function  $V$  (cf. Section III).

Note that different  $\delta$ -ISS- $M_{\mathbf{q}}$  Lyapunov functions  $V$  result in different quantization parameters  $\mathbf{q}$  satisfying (V.3) and (V.4). Hence, one can seek appropriate  $\delta$ -ISS- $M_{\mathbf{q}}$  Lyapunov functions to provide less conservative quantization parameters  $\mathbf{q}$  (cf. Remark 3.6). For example, one can leverage the existing techniques in the literature such as [23] to construct polyhedral incremental Lyapunov functions which are less conservative compared to quadratic ones.

Note that the results in [16], as in the following corollary, are fully recovered by the statement in Theorem 5.1 if the stochastic control system  $\Sigma$  is not affected by any noise,

implying that  $h(\sigma, \tau)$  is identically zero and that the  $\delta$ -ISS- $M_q$  property reduces to the  $\delta$ -ISS property.

*Corollary 5.2:* Let  $\bar{\Sigma}$ , the non-probabilistic version of  $\Sigma$ , be a  $\delta$ -ISS control system admitting a  $\delta$ -ISS Lyapunov function  $V$ . For any  $\varepsilon \in \mathbb{R}^+$ , and any triple  $\mathbf{q} = (\tau, \eta, \mu)$  of quantization parameters satisfying  $\mu \leq \text{span}(\mathbf{U})$  and

$$\bar{\alpha}(\eta) \leq \underline{\alpha}(\varepsilon), \quad (\text{V.6})$$

$$e^{-\kappa\tau} \underline{\alpha}(\varepsilon) + \frac{1}{e^{\kappa}} \rho(\mu) + \hat{\gamma}(\eta) \leq \underline{\alpha}(\varepsilon), \quad (\text{V.7})$$

one obtains  $S_{\mathbf{q}}(\Sigma) \cong_{\mathcal{S}}^{\varepsilon} S_{\tau}(\bar{\Sigma})$ .

Note that since any non-probabilistic point can be regarded as a random variable with a Dirac probability distribution centered at that point, the output set of  $S_{\tau}(\bar{\Sigma})$ , which is  $\mathbb{R}^n$ , can be embedded in the output set of  $S_{\mathbf{q}}(\Sigma)$ . Hence, without loss of generality, one can assume that the output sets of  $S_{\tau}(\bar{\Sigma})$  and  $S_{\mathbf{q}}(\Sigma)$  are equal. The same remark holds for Corollary 5.4.

The next main theorem provides a result that is similar to that in Theorem 5.1, which is however not obtained by explicit use of the parameters of  $\delta$ -ISS- $M_q$  Lyapunov functions (e.g.  $\underline{\alpha}$ ,  $\bar{\alpha}$ ,  $\kappa$ ,  $\rho$ ,  $\hat{\gamma}$ ), but by using functions  $\beta$  and  $\gamma$  as in (III.1). Note that the symbolic models computed by using the quantization parameters  $\mathbf{q}$  provided in the next theorem, whenever existing, is likely to have fewer states than the ones computed by using the quantization parameters  $\mathbf{q}$  provided in Theorem 5.1 based on our experiences in different simulations and the non-probabilistic version of this theorem as recalled in Corollary 5.4. It also may provide a better lower bound on  $\varepsilon$  for a fixed sampling time  $\tau$ . We refer the interested reader to the case studies section for a comparison between the results of Theorem 5.1 and of the next theorem on some practical examples.

*Theorem 5.3:* Consider a  $\delta$ -ISS- $M_q$  stochastic control system  $\Sigma$ , satisfying (III.8). For any  $\varepsilon \in \mathbb{R}^+$ , and any triple  $\mathbf{q} = (\tau, \eta, \mu)$  of quantization parameters satisfying  $\mu \leq \text{span}(\mathbf{U})$  and

$$(\beta(\varepsilon^q, \tau) + \gamma(\mu))^{\frac{1}{q}} + (h(\sigma, \tau))^{\frac{1}{q}} + \eta \leq \varepsilon, \quad (\text{V.8})$$

we have  $S_{\mathbf{q}}(\Sigma) \cong_{\mathcal{S}}^{\varepsilon} S_{\tau}(\Sigma)$ .

The proof of Theorem 5.3 is provided in the Appendix.

It can be readily seen that when we are interested in the dynamics of  $\Sigma$ , initialized on a compact  $D \subset \mathbb{R}^n$  of the form of finite union of boxes and for a given precision  $\varepsilon$ , there always exists a sufficiently large value of  $\tau$  and small values of  $\eta$  and  $\mu$  such that  $\eta \leq \text{span}(D)$  and the condition in (V.8) are satisfied. However, unlike the result in Theorem 5.1, notice that here for a given fixed sampling time  $\tau$ , one may not find any values of  $\eta$  and  $\mu$  satisfying (V.8) because the quantity  $(\beta(\varepsilon^q, \tau))^{\frac{1}{q}}$  may be larger than  $\varepsilon$ . As long as there exists a triple  $\mathbf{q}$ , satisfying (V.8), the lower bound of precision  $\varepsilon$  can be computed by solving the following inequality with respect to  $\varepsilon$ :  $\varepsilon - (\beta(\varepsilon^q, \tau))^{\frac{1}{q}} > (h(\sigma, \tau))^{\frac{1}{q}}$ . In this case, one can easily verify that the lower bound on  $\varepsilon$  goes to zero as  $\tau$  goes to infinity, as  $\tau$  goes to zero (only if  $(\beta(\varepsilon^q, 0))^{\frac{1}{q}} \leq \varepsilon$ ), or as  $Z \rightarrow 0$ , where  $Z$  is the Lipschitz constant introduced in Definition 2.1.

Note that the results in [34], as in the following corollary, are fully recovered by the results in Theorem 5.3 if the stochastic control system  $\Sigma$  is not affected by any noise, implying that  $h(\sigma, \tau)$  is identically zero and that the  $\delta$ -ISS- $M_q$

property becomes the  $\delta$ -ISS property.

*Corollary 5.4:* Let  $\bar{\Sigma}$ , the corresponding non-probabilistic version of  $\Sigma$ , be a  $\delta$ -ISS control system (i.e. satisfying (III.2)). For any  $\varepsilon \in \mathbb{R}^+$ , and any triple  $\mathbf{q} = (\tau, \eta, \mu)$  of quantization parameters satisfying  $\mu \leq \text{span}(\mathbf{U})$  and

$$\beta(\varepsilon, \tau) + \gamma(\mu) + \eta \leq \varepsilon, \quad (\text{V.9})$$

we have  $S_{\mathbf{q}}(\Sigma) \cong_{\mathcal{S}}^{\varepsilon} S_{\tau}(\bar{\Sigma})$ .

Based upon the proposed symbolic models in Theorems 5.1 and 5.3, one can leverage the automata-theoretic controller synthesis algorithms [26] to synthesize hybrid controllers enforcing complex logic specifications (e.g. LTL) on the original stochastic control systems. Particularly, we refer the interested readers to Proposition 11.10 in [39] elucidating how a controller synthesized to solve a simulation game over a symbolic model can be refined to a controller for the original system; for detailed definitions of syntax and semantics of LTL, see also [6, Chapter 5].

Although  $S_{\tau}(\Sigma)$  inherits a classical trace-based semantics [39], the outputs of  $S_{\tau}(\Sigma)$  being random variables makes it less trivial especially in terms of satisfaction of temporal logic specifications, e.g. LTL. Let us focus on the following example. Assume  $W \subset \mathbb{R}^n$  to be the safe set of our interest, i.e. the LTL formula  $\Box W$ . Suppose we are able to find a control strategy over the abstraction  $S_{\mathbf{q}}(\Sigma)$  enforcing an output run of  $S_{\mathbf{q}}(\Sigma)$  to satisfy  $\Box W$ . Since the output run consists of points  $y \in Y_{\mathbf{q}} = Y_{\tau}$  such that  $y \in W \subset \mathbb{R}^n$ ,  $y$  is a degenerate random variable. On the other hand, assume that  $S_{\tau}(\Sigma) \cong_{\mathcal{S}}^{\varepsilon} S_{\mathbf{q}}(\Sigma)$ . By interpreting the result  $\Box W$  obtained over the abstraction, one can guarantee that the corresponding output run of  $S_{\tau}(\Sigma)$  satisfies  $\Box W_{\varepsilon}$ , where  $W_{\varepsilon} = \{y \in Y_{\tau} \mid \mathbf{d}(y, W) \leq \varepsilon\}$ ,  $\mathbf{d}(y, W) = \inf_{w \in W} \mathbf{d}(y, w)$ , and  $\mathbf{d}$  is the  $q$ th moment metric. Note that although the original set  $W$  is a subset of  $\mathbb{R}^n$ , its  $\varepsilon$ -inflation in the  $q$ th moment metric, i.e.  $W_{\varepsilon}$ , is not a subset of  $\mathbb{R}^n$  anymore and contains non-degenerate random variables. As a result, although satisfying  $\Box W_{\varepsilon}$  does not necessarily mean that a trajectory of  $\Sigma$  always stays within some non-probabilistic set, it means that the associated random variables always belong to  $W_{\varepsilon}$  and hence are  $\varepsilon$ -close to the non-probabilistic set  $W$  with respect to the  $q$ th moment metric.

*Remark 5.5:* Although we assume that the set  $\mathbf{U}$  is infinite, Theorems 5.1 (resp. Corollary 5.2) and 5.3 (resp. Corollary 5.4) still hold when the set  $\mathbf{U}$  is finite, with the following modifications. First, the system  $\Sigma$  (resp.  $\bar{\Sigma}$ ) is required to satisfy the property (III.1) (resp. (III.2)) for  $v = v'$ . Second, take  $U_{\mathbf{q}} = \mathbf{U}$  in the definition of  $S_{\mathbf{q}}(\Sigma)$ . Finally, in the conditions (V.4) (resp. (V.7)) and (V.8) (resp. (V.9)) set  $\mu = 0$ .  $\square$

If the stochastic control system  $\Sigma$  is not  $\delta$ -ISS- $M_q$ , one can use the results in [44], providing symbolic models that are only sufficient in the sense that the refinement of any controller synthesized for the symbolic model enforces the same desired specifications on the original stochastic control system (in the sense of moments). However, they can no longer guarantee, as it was the case in this paper, that the existence of a controller for the original stochastic control system leads to the existence of a controller for the symbolic model.  $\square$

*Remark 5.6:* A detailed computational complexity analysis of symbolic controller synthesis using the proposed abstractions is provided in Section V in [33]. In particular, both Algorithms 1 and 2 in [33] can be applied to synthesize symbolic

controllers based upon the proposed symbolic abstractions in this paper.

### B. Relationship with notions of probabilistic bisimulation in the literature

In the remainder of this section, we relate the (approximate) equivalence relations introduced in this work with known probabilistic concepts in the literature.

The approximate bisimulation notion in Definition 4.2 is structurally different from the probabilistic version discussed for finite state, *discrete-time* labeled Markov chains in [13], which is also extended to continuous-space processes as in [12] and hinges on the absolute difference between transition probabilities over sets covering the state space. The notion in this work can be instead related to the approximate probabilistic bisimulation notion discussed in [19], which lower bounds the probability that the Euclidean distance between any trace of the abstract model and the corresponding one of the concrete model and vice versa remains close: both notions hinge on distances over trajectories, rather than over transition probabilities as in [12], [13].

We make the above statement more precise with the following results, which do not require the assumptions  $f(0_n, 0_m) = 0_n$  and  $\sigma(0_n) = 0_{n \times p}$  anymore.

**Proposition 5.7:** Let  $\Sigma$  be a stochastic control system. Assume that there exists a stochastic bisimulation function  $\phi : \mathbb{R}^n \times \mathbb{R}^n \rightarrow \mathbb{R}_0^+$ , as defined in [19], between  $\Sigma$  and its corresponding non-probabilistic control system  $\bar{\Sigma}$ , and assume that  $\bar{\Sigma}$  is  $\delta$ -ISS. For any  $\varepsilon \in \mathbb{R}^+$ , any triple  $\mathbf{q} = (\tau, \eta, \mu)$  of quantization parameters, satisfying  $\mu \leq \text{span}(\mathbf{U})$  and (V.9) (or (V.6) and (V.7)), and for any state run  $\xi_{x_{\tau_0} v}$  of  $S_\tau(\Sigma)$ , there exists a state run  $X_{x_{q_0} u}$  of  $S_q(\Sigma)$ , and vice versa, such that the following relation holds:

$$\mathbb{P} \left\{ \sup_{k \in \mathbb{N}_0} \|\xi_{x_{\tau_0} v}(k\tau) - X_{x_{q_0} u}(k)\| > \varepsilon \mid x_{\tau_0} \right\} \leq \frac{\sqrt{\phi(x_{\tau_0}, x_{\tau_0})} + \varepsilon}{\varepsilon}. \quad (\text{V.10})$$

The proof of Proposition 5.7 is provided in the Appendix.

Theorem 3 in [19] provides a similar result as in Proposition 5.7, however it is limited to hold over two infinite systems rather than a finite and an infinite one as in this work. As explained in [19], in order to compute a stochastic bisimulation function, one requires to solve a game, which is computationally difficult even for linear stochastic control systems. On the other hand, if the set of input is a singleton (i.e.,  $\mathbf{U} = \{0_m\}$ ), which results in the verification of stochastic dynamical systems, then a stochastic bisimulation function  $\phi$  can be efficiently computed by solving some LMIs, as explained in [19, equations (39) and (40)] for linear stochastic control systems. Alternatively, the stochastic contractivity of the model can be used for this goal [1].

**Remark 5.8:** Let us further comment on the relationship between  $\delta$ -ISS- $M_q$  Lyapunov functions in this work and stochastic bisimulation ones in [19]. Let  $\Sigma = (\mathbb{R}^n, \{0_m\}, \mathcal{U}_\tau, f, \sigma)$  be a stochastic control system admitting a  $\delta$ -ISS- $M_q$  Lyapunov function  $V$ , and such that  $f(0_n, 0_m) = 0_n$ ,  $\sigma(0_n) = 0_{n \times p}$ , and  $0_{n \times n} \preceq \partial_{\bar{x}, \bar{x}} V(0_n, \bar{x})$  for all  $\bar{x} \in \mathbb{R}^n$ . It can be readily verified that the function  $\phi(x, \bar{x}) = V(x, 0_n) + V(0_n, \bar{x})$ , for all  $x, \bar{x} \in \mathbb{R}^n$ , is a stochastic bisimulation function between  $\Sigma$  and

the corresponding non-probabilistic control system  $\bar{\Sigma}$ , under the following modification in condition (i) in [19, Definition 2]: (i)  $\phi(x, \bar{x}) \geq \tilde{\alpha}(\|x - \bar{x}\|)$ , for some convex  $\tilde{\alpha} \in \mathcal{K}_\infty$  and all  $x, \bar{x} \in \mathbb{R}^n$ . Using this new condition on  $\phi$ , one can readily revise the relation (V.10) correspondingly.  $\square$

The next theorem, which is computationally more tractable for stochastic control systems and does not require the existence of stochastic bisimulation functions as in [19], provides a similar result as in Proposition 5.7, but applies only over a finite time horizon.

**Proposition 5.9:** Let  $\Sigma$  be a stochastic control system. Suppose there exist a  $\delta$ -ISS- $M_q$  Lyapunov function  $V$  for  $\Sigma$  and  $\alpha \in \mathbb{R}_0^+$  such that

$$\text{Tr}(\sigma(x)^T \partial_{x,x} V(x, y) \sigma(x)) - \text{Tr} \left( \begin{bmatrix} \sigma(x) \\ \sigma(y) \end{bmatrix} \begin{bmatrix} \sigma^T(x) & \sigma^T(y) \end{bmatrix} H(V)(x, y) \right) \leq \alpha,$$

for all  $x, y \in \mathbb{R}^n$ . For any  $\varepsilon \in \mathbb{R}^+$ , any triple  $\mathbf{q} = (\tau, \eta, \mu)$  of quantization parameters, satisfying  $\mu \leq \text{span}(\mathbf{U})$  and (V.9) (or (V.6), and (V.7)), and for any state run  $\xi_{x_{\tau_0} v}$  of  $S_\tau(\Sigma)$ , there exists a state run  $X_{x_{q_0} u}$  of  $S_q(\Sigma)$ , and vice versa, such that the following relations hold:

$$\mathbb{P} \left\{ \sup_{0 \leq k \leq N} \|\xi_{x_{\tau_0} v}(k\tau) - X_{x_{q_0} u}(k)\| > \varepsilon + \varepsilon \mid x_{\tau_0} \right\} \leq 1 - e^{-\frac{\alpha N \tau}{2\alpha(\varepsilon^q)}}, \text{ if } \underline{\alpha}(\varepsilon^q) \geq \frac{\alpha}{2\kappa}, \quad (\text{V.11})$$

$$\mathbb{P} \left\{ \sup_{0 \leq k \leq N} \|\xi_{x_{\tau_0} v}(k\tau) - X_{x_{q_0} u}(k)\| > \varepsilon + \varepsilon \mid x_{\tau_0} \right\} \leq \frac{(e^{N\tau\kappa} - 1)\alpha}{2\kappa\underline{\alpha}(\varepsilon^q) e^{N\tau\kappa}}, \text{ if } \underline{\alpha}(\varepsilon^q) \leq \frac{\alpha}{2\kappa}. \quad (\text{V.12})$$

The proof of Proposition 5.9 is provided in the Appendix.

As an alternative to the two results above, we now introduce a probabilistic approximate bisimulation relation between  $S_\tau(\Sigma)$  and  $S_q(\Sigma)$  point-wise in time: this relation is sufficient to work with LTL specifications for which satisfiability can be ascertained at single time instances, such as for next ( $\circ$ ) and eventually ( $\diamond$ ). Please look at the explanation after the proof of Proposition 5.11 for more details.

**Definition 5.10:** Consider two systems  $S_\tau(\Sigma)$  and  $S_q(\Sigma)$ , and precisions  $\varepsilon \in \mathbb{R}^+$  and  $\hat{\varepsilon} \in [0, 1]$ . A relation  $R \subseteq X_\tau \times X_q$  is said to be an  $(\varepsilon, \hat{\varepsilon})$ -approximate simulation relation from  $S_\tau$  to  $S_q$ , if the following three conditions are satisfied:

- (i) for every  $x_{\tau_0} \in X_{\tau_0}$ , there exists  $x_{q_0} \in X_{q_0}$  with  $(x_{\tau_0}, x_{q_0}) \in R$ ;
- (ii) for every  $(x_\tau, x_q) \in R$ , we have  $\mathbb{P} \{ \|H_\tau(x_\tau) - H_q(x_q)\| \geq \varepsilon \} \leq \hat{\varepsilon}$ ;
- (iii) for every  $(x_\tau, x_q) \in R$ , we have that  $x_\tau \xrightarrow{v_\tau} x'_\tau$  in  $S_\tau$  implies the existence of  $x_q \xrightarrow{u_q} x'_q$  in  $S_q$  satisfying  $(x'_\tau, x'_q) \in R$ .

A relation  $R \subseteq X_\tau \times X_q$  is said to be an  $(\varepsilon, \hat{\varepsilon})$ -approximate bisimulation relation between  $S_\tau$  and  $S_q$  if  $R$  is an  $(\varepsilon, \hat{\varepsilon})$ -approximate simulation relation from  $S_\tau$  to  $S_q$  and  $R^{-1}$  is an  $(\varepsilon, \hat{\varepsilon})$ -approximate simulation relation from  $S_q$  to  $S_\tau$ .

System  $S_\tau$  is  $(\varepsilon, \hat{\varepsilon})$ -approximately simulated by  $S_q$ , or  $S_q$   $(\varepsilon, \hat{\varepsilon})$ -approximately simulates  $S_\tau$ , denoted by  $S_\tau \preceq_S^{(\varepsilon, \hat{\varepsilon})} S_q$ , if there exists an  $(\varepsilon, \hat{\varepsilon})$ -approximate simulation relation from  $S_\tau$  to  $S_q$ . System  $S_\tau$  is  $(\varepsilon, \hat{\varepsilon})$ -approximate bisimilar to  $S_q$ ,



denoted by  $S_\tau \cong_S^{(\epsilon, \hat{\epsilon})} S_q$ , if there exists an  $(\epsilon, \hat{\epsilon})$ -approximate bisimulation relation between  $S_\tau$  and  $S_q$ .

We show next the existence of symbolic models with respect to the notion of approximate bisimulation relation in Definition 5.10.

**Proposition 5.11:** Let  $\Sigma$  be a stochastic control system. For any  $\epsilon \in \mathbb{R}^+$  and  $\hat{\epsilon} \in [0, 1]$ , we have  $S_\tau(\Sigma) \cong_S^{(\epsilon, \hat{\epsilon})} S_q(\Sigma)$  if  $S_\tau(\Sigma) \cong_S^\epsilon S_q(\Sigma)$  for  $\epsilon = \hat{\epsilon}$ .

*Proof:* The proof is a simple consequence of Theorems 5.1 or 5.3 and Markov inequality [30], used as the following:

$$\begin{aligned} \mathbb{P} \{ \|H_\tau(x_\tau) - H_q(x_q)\| \geq \epsilon \} &\leq \frac{\mathbb{E} \{ \|H_\tau(x_\tau) - H_q(x_q)\| \}}{\epsilon} \\ &\leq \frac{(\mathbb{E} \{ \|H_\tau(x_\tau) - H_q(x_q)\|^q \})^{\frac{1}{q}}}{\epsilon} \leq \frac{\epsilon}{\epsilon} = \hat{\epsilon}. \end{aligned}$$

Let us clarify the use of the previous proposition in a simple example. Given a stochastic control system  $\Sigma$ , assume there exists a symbolic model  $S_q(\Sigma)$  such that  $S_\tau(\Sigma) \cong_S^\epsilon S_q(\Sigma)$ . If there exists an output run  $X_{x_0 u}$  of  $S_q(\Sigma)$  reaching a set  $W$ , i.e. satisfying the LTL formula  $\diamond W$ , then for the corresponding output run  $\xi_{x_0 v}$  of  $S_\tau(\Sigma)$ , one has

$$\begin{aligned} \mathbb{P} \{ \xi_{x_0 v} \models \diamond W^\epsilon \} &:= \mathbb{P} \{ \exists N \in \mathbb{N} \text{ s.t. } \xi_{x_0 v}(N\tau) \in W^\epsilon \} \\ &\geq \sup_{M \in \mathbb{N}} \mathbb{P} \{ \xi_{x_0 v}(M\tau) \in W^\epsilon \} \geq 1 - \frac{\epsilon}{\epsilon}, \end{aligned} \quad (\text{V.13})$$

where the last inequality results from Proposition 5.11. Hence, one can leverage the result of Proposition 5.11 to provide a lower bound on the probability of satisfaction of logic specifications for which satisfiability can be obtained at single time instances. Note that this is not the case for logic specifications including the (bounded) quantifier ‘‘ $\forall$ ’’ such as always ( $\square$ ) or even until ( $\mathcal{U}$ ). By taking out the quantifier  $\forall$  from the probability operator, the first inequality sign in (V.13) reverses and hence we cannot provide the lower bound on the probability of satisfaction anymore.

## VI. CASE STUDIES

We now experimentally demonstrate the effectiveness of the discussed results. In the examples below, the computation of the abstract systems  $S_q(\Sigma)$  have been implemented by the software tool *Pessoa* [27] on a laptop with CPU 2GHz Intel Core i7. We have assumed that the control inputs are piecewise constant of duration  $\tau$  and that  $\mathcal{U}_\tau$  is finite and contains signals taking values in  $[U]_\mu$ . Hence, as explained in Remark 5.5,  $\mu = 0$  is to be used in the conditions (V.4) and (V.8). The controllers enforcing the specifications of interest have been found by standard algorithms from game theory [26], as implemented in *Pessoa*. In both examples, the terms  $W_t^i, i = 1, 2$ , denote the standard Brownian motion terms. Although we used *Pessoa* to construct bisimilar symbolic models for the examples below, one can use the constructed symbolic models as an input for other existing synthesis toolboxes such as SPIN [18] to synthesize symbolic controllers enforcing larger classes of LTL specifications rather than the limited ones supported by *Pessoa*.

### A. Nonlinear model, 2nd moment, ‘‘sequential target tracking’’ property

Consider the nonlinear model of a pendulum on a cart borrowed from [9], which is now affected by noise. The model

is described by:

$$\Sigma : \begin{cases} d\xi_1 = \xi_2 dt + 0.03\xi_1 dW_t^1, \\ d\xi_2 = \left(-\frac{g}{l} \sin(\xi_1) - \frac{k}{m}\xi_2 + \frac{1}{ml^2}v\right) dt + 0.03\xi_2 dW_t^1, \end{cases} \quad (\text{VI.1})$$

where  $\xi_1$  and  $\xi_2$  represent the angular position and the velocity of the point mass on the pendulum,  $v$  is the torque applied to the cart,  $g = 9.8$  is acceleration due to gravity,  $l = 0.5$  is the length of the shaft,  $m = 0.6$  is the mass, and  $k = 2$  is the friction coefficient. All the constants and the variables are considered in SI units. We assume that  $U = [-1.5 \ 1.5]$  and that  $\mathcal{U}_\tau$  contains signals taking values in  $[U]_{0.5}$ . We work on the subset  $D = [-1 \ 1] \times [-1 \ 1]$  of the state space of  $\Sigma$ . Using the Lyapunov function  $V(x, x') = (x - x')^T P (x - x')$ , for all  $x, x' \in \mathbb{R}^2$ , proposed in [9], where  $P = [1.5 \ 0.3; 0.3 \ 1.5]$ , it can be readily verified that  $V$  satisfies conditions (i) and (ii) in Definition 3.2 with  $\underline{\alpha}(r) = 1.2r$ , and  $\bar{\alpha}(r) = 3.6r$ , for  $q = 2$ . Moreover,  $V$  satisfies condition (iii) in Definition 3.2 with  $\kappa = 0.7691$ , and  $\rho(r) = 8.76r$ , for all  $x, x' \in D$ , and again  $q = 2$ . Therefore,  $\Sigma$  is  $\delta$ -ISS- $M_2$ , equipped with the  $\delta$ -ISS- $M_2$  Lyapunov function  $V$ , as long as we are interested in its dynamics, initialized on  $D$ . Using the results in Theorem 3.3, one obtains that functions  $\beta(r, s) = 3e^{-\kappa s}$  and  $\gamma(r) = 3.49r$  satisfy property (III.1) for  $\Sigma$ .

For a given fixed sampling time  $\tau = 3$ , the precision  $\epsilon$  is lower bounded by 0.0778 and 0.4848 using the results in Theorems 5.3 and 5.1, respectively. Hence, the results in Theorem 5.3 provide a symbolic model which is much less conservative than the one provided by Theorem 5.1. By selecting  $\epsilon = 0.085$ , the parameter  $\eta$  for  $S_q(\Sigma)$  based on the results in Theorem 5.3 is obtained as 0.0033. The resulting cardinality of the state and input sets for  $S_q(\Sigma)$  are 370881 and 7, respectively. The CPU time taken for computing the abstraction  $S_q(\Sigma)$  has amounted to 17278 seconds.

Now, consider the objective to design a controller forcing the trajectories of  $\Sigma$ , starting from the initial condition  $x_0 = (0, 0)$ , to first sequentially visit two regions of interest  $W_1 = [0.48 \ 0.58] \times [-1 \ 1]$ , and  $W_2 = [-0.58 \ -0.48] \times [-1 \ 1]$ ; then, once the system has visited the regions, to return to the first region  $W_1$  and remain there forever. The LTL formula<sup>1</sup> encoding this goal is  $\diamond \square W_1 \wedge \diamond (W_1 \wedge \diamond W_2)$ . The CPU time taken for synthesizing the controller has amounted to 3.02 seconds. Figure 1 displays a few realizations of the closed-loop solution process  $\xi_{x_0 v}$  stemming from the initial condition  $x_0 = (0, 0)$ , as well as the corresponding evolution of the input signal. In Figure 2, we show the square root of the average value (over 100 experiments) of the squared distance in time of the solution process  $\xi_{x_0 v}$  to the sets  $W_1$  and  $W_2$ , namely,  $\|\xi_{x_0 v}(t)\|_{W_1}^2$  and  $\|\xi_{x_0 v}(t)\|_{W_2}^2$ . Notice that the square root of this empirical (averaged) squared distances is significantly lower than the selected precision  $\epsilon = 0.085$ , as expected since the conditions based on Lyapunov functions can lead to conservative bounds. (We have discussed that the bounds can be improved by seeking optimized Lyapunov functions.)

Notice that using the chosen quantization parameters and (V.9), we obtain a precision  $\epsilon = 0.005$  to be used in (V.11) or (V.12). As long as we are interested in dynamics of  $\Sigma$  on  $D$ , we have an  $\alpha = 0.0032$  in Proposition 5.9. Using the result

<sup>1</sup>Note that the semantics of LTL are defined over the output behaviors of  $S_q(\Sigma)$ .

in Proposition 5.9, one can conclude that the refinement of a control policy satisfying a bounded horizon LTL (BLTL) formula<sup>2</sup>  $\varphi$  over a discrete time horizon spanning  $\{0, 3, \dots, 27\}$  seconds for  $S_q(\Sigma)$  (e.g.  $\varphi = \diamond \square W_1 \wedge \diamond (W_1 \wedge \diamond W_2)$ ) to the system  $S_\tau(\Sigma)$  satisfies the “inflated formula”<sup>3</sup>  $\varphi^{0.31}$  (e.g.  $\varphi^{0.31} = \diamond \square W_1^{0.31} \wedge \diamond (W_1^{0.31} \wedge \diamond W_2^{0.31})$ ) with probability at least 70%. To get the better lower bound for the aforementioned probability, one can reduce the time horizon, increase the inflation factor, or do both of them simultaneously. For example, by considering the discrete time horizon  $\{0, 3, \dots, 15\}$  rather than  $\{0, 3, \dots, 27\}$  and inflation factor 0.5 rather than 0.31, the lower bound of the previous probability will be 92%. The same observation works for the next example as well.

Furthermore, employing the result in Proposition 5.11, one can conclude that the refinement of a controller satisfying  $\diamond W$  for  $S_q(\Sigma)$  to system  $S_\tau(\Sigma)$  satisfies  $\diamond W^\epsilon$  with probability at least  $1 - \epsilon/\epsilon$ . For example, refinement of a controller, satisfying  $\diamond W_1$  for  $S_q(\Sigma)$ , to the system  $S_\tau(\Sigma)$  satisfies  $\diamond W_1^{0.28}$  with probability at least 70%. By changing the inflation factor from 0.28 to 0.6, the lower bound of the previous probability will be 0.86%.




Fig. 1. Example 1: A few realizations of the closed-loop solution process  $\xi_{x_0 v}$  with initial condition  $x_0 = (0, 0)$  (bottom panel) and the evolution of the input signal  $v$  (top panel).




Fig. 2. Example 1: Square root of the average values (over 100 experiments) of the squared distance of the solution process  $\xi_{x_0 v}$  to the sets  $W_1$  (left panels) and  $W_2$  (right panels), in two different vertical scales (top vs bottom panels).

<sup>2</sup>To compute the probability of satisfaction of logic specifications based on the result of Proposition 5.9, we need to restrict ourselves to the bounded horizon specifications (e.g. BLTL). We refer the readers to [8] for the detailed definition of bounded LTL (BLTL).

<sup>3</sup>We refer the interested readers to [24, Definition 1] for a detailed definition of the inflated version of a BLTL formula.

## B. Linear model, 1st moment, “reach-and-stay, while staying” property

Consider a linear DC motor borrowed from [28], now affected by noise and described by:

$$\Sigma : \begin{cases} d\xi_1 = \left(-\frac{b}{J}\xi_1 + \frac{K}{J}\xi_2\right) dt + 0.15\xi_1 dW_t^1, \\ d\xi_2 = \left(-\frac{K}{L}\xi_1 - \frac{R}{L}\xi_2 + \frac{1}{L}v\right) dt + 0.15\xi_2 dW_t^2, \end{cases} \quad (\text{VI.2})$$

where  $\xi_1$  is the angular velocity of the motor,  $\xi_2$  is the current through an inductor,  $v$  is the voltage signal,  $b = 10^{-4}$  is the damping ratio of the mechanical system,  $J = 25 \times 10^{-5}$  is the moment of inertia of the rotor,  $K = 5 \times 10^{-2}$  is the electromotive force constant,  $L = 3 \times 10^{-4}$  is the electric inductance, and  $R = 0.5$  is the electric resistance. All constants and variables are considered in the SI units. It can be readily verified that the system  $\Sigma$  satisfies condition (III.6) with constant  $\hat{\kappa} = 40$  and matrix  $P = [1.2201 \ 0.2224; 0.2224 \ 1.2248]$ . Therefore,  $\Sigma$  is  $\delta$ -ISS- $M_q$  and equipped with the  $\delta$ -ISS- $M_q$  Lyapunov function  $V(x, x')$  in (III.3), where  $q \in \{1, 2\}$ . In this example, we use  $q = 1$ .

We assume that  $U = [-0.5 \ 0.5]$  and that  $\mathcal{U}_\tau$  contains signals taking values in  $[U]_{0.1}$ . We work on the subset  $D = [-5 \ 5] \times [-5 \ 5]$  of the state space of  $\Sigma$ . For a given precision  $\epsilon = 1$  and fixed sampling time  $\tau = 0.01$ , the parameter  $\eta$  of  $S_q(\Sigma)$  based on the results in Theorem 5.1 is equal to 0.01. Note that for sampling times  $\tau < 0.026$ , and in particular for a choice  $\tau = 0.01$ , the results in Theorem 5.3 cannot be applied here because  $(\beta(\epsilon^q, \tau))^{\frac{1}{q}} > \epsilon$  and condition (V.8) in Theorem 5.3 is not fulfilled. The resulting cardinality of the state and input sets for  $S_q(\Sigma)$  amounts to 1002001 and 11, respectively. The CPU time used for computing the abstraction has amounted to 148.092 seconds.

Now, consider the objective to design a controller forcing the trajectories of  $\Sigma$  to reach and stay within  $W = [4.5 \ 5] \times [-0.25 \ 0.25]$ , while ensuring that the current through the inductor is restricted between  $-0.25$  and  $0.25$ . This corresponds to the LTL specification  $\diamond \square W \wedge \square Z$ , where  $Z = [-5 \ 5] \times [-0.25 \ 0.25]$ . The CPU time used for synthesizing the controller has been of 3.88 seconds. Figure 3 displays a few realizations of the closed-loop solution process  $\xi_{x_0 v}$  stemming from the initial condition  $x_0 = (-5, -0)$ , as well as the corresponding evolution of the input signal. In Figure 4, we show the average value (over 100 experiments) of the distance in time of the solution process  $\xi_{x_0 v}$  to the sets  $W$  and  $Z$ , namely  $\|\xi_{x_0 v}(t)\|_W$  and  $\|\xi_{x_0 v}(t)\|_Z$ . Notice that the empirical average distances are as expected significantly lower than the precision  $\epsilon = 1$ .

Note that using the selected quantization parameters in (V.6) and (V.7), we obtain a value  $\epsilon = 0.08$  in (V.10), (V.11) or (V.12). Further, using the result in Proposition 5.7, one obtains that, as long as we are interested in dynamics of  $\Sigma$ , initialized on  $D$  and  $U = \{0_m\}$ , there exists a state run  $X_{x_0 u}$  of  $S_q(\Sigma)$  satisfying an LTL formula  $\varphi$  if and only if there exists a state run  $\xi_{x_{\tau_0} v}$  of  $S_\tau(\Sigma)$  satisfying  $\varphi^1$  with probability at least 79%, where  $\varphi^1$  is the 1-inflation of  $\varphi$ .

Moreover, using the result in Proposition 5.9, one can conclude that, as long as we are interested in dynamics of  $\Sigma$  on  $Z$ , implying that  $\alpha = 0.6917$  in Proposition 5.9, the refinement of a controller, satisfying a BLTL formula  $\varphi$  for  $S_q(\Sigma)$  (e.g.  $\varphi = \diamond \square W \wedge \square Z$ ) to the system  $S_\tau(\Sigma)$  satisfies  $\varphi^1$  (e.g.  $\varphi^1 = \diamond \square W^1 \wedge \square Z^1$ ) with probability at least 70%

over a discrete time horizon spanning  $\{0, 0.01, 0.02, \dots, 1\}$  seconds.




Fig. 3. Example 2: A few realizations of the closed-loop solution process  $\xi_{x_0 v}$  with initial condition  $x_0 = (-5, 0)$  in different scales (left and middle panel), and the evolution of the input signal  $v$  (right panel).




Fig. 4. Example 2: Average values (over 100 experiments) of the distance of the solution process  $\xi_{x_0 v}$  to the sets  $W$  (left panels) and  $Z$  (right panels), in two different vertical scales (top vs bottom panels).

## VII. CONCLUSIONS

This work has shown that any stochastic sampled-data control system, admitting a  $\delta$ -ISS- $M_q$  Lyapunov function, and initializing within a compact set of states, admits a finite approximately bisimilar symbolic model (in the sense of moments or probability). The constructed symbolic model can be used to synthesize controllers enforcing complex logic specifications, expressed via linear temporal logic or as automata on infinite strings.

The main limitation of the design methodology developed in this paper lies in the cardinality of the set of states of the computed symbolic model. The authors are currently investigating several different techniques to address this limitation. For example, the recent work in [47] by some of the authors, based upon the results of this article, provides a new way of constructing bisimilar symbolic abstractions which does not require discretization of the set of states but only the set of inputs. Hence, it can be potentially more tractable.

## VIII. ACKNOWLEDGEMENTS

The authors would like to thank Ilya Tkachev for fruitful technical discussions.

## REFERENCES

- [1] A. Abate. A contractivity approach for probabilistic bisimulations of diffusion processes. In *Proceedings of the 48th IEEE Conference on Decision and Control*, pages 2230–2235, December 2009.

- [2] A. Abate, S. Amin, M. Prandini, J. Lygeros, and S. Sastry. Computational approaches to reachability analysis of stochastic hybrid systems. In *Proceedings of the 10th International Conference on Hybrid Systems: Computation and Control*, pages 4–17, April 2007.
- [3] A. Abate, A. D’Innocenzo, and M. D. Di Benedetto. Approximate abstractions of stochastic hybrid systems. *IEEE Transactions on Automatic Control*, 56(11):2688–2694, November 2011.
- [4] D. Angeli. A Lyapunov approach to incremental stability properties. *IEEE Transactions on Automatic Control*, 47(3):410–21, March 2002.
- [5] S. I. Azuma and G. J. Pappas. Discrete abstraction of stochastic nonlinear systems: a bisimulation function approach. In *Proceedings of American Control Conference*, pages 1035–1040, June 2010.
- [6] C. Baier and J. P. Katoen. *Principles of model checking*. The MIT Press, April 2008.
- [7] C. Belta and L. C. G. J. M. Habets. Controlling a class of nonlinear systems on rectangles. *IEEE Transactions on Automatic Control*, 51(11):1749–1759, November 2006.
- [8] A. Biere, A. Cimatti, E. M. Clarke, and Y. Zhu. Symbolic model checking without bdds. In *Proceedings of the 5th International Conference on Tools and Algorithms for Construction and Analysis of Systems*, pages 193–207, March 1999.
- [9] A. Borri, G. Pola, and M. D. Di Benedetto. Symbolic models for nonlinear control systems affected by disturbances. *International Journal of Control*, 85(10):1422–1432, May 2012.
- [10] D. Chatterjee and D. Liberzon. Stability analysis of deterministic and stochastic switched systems via a comparison principle and multiple Lyapunov functions. *SIAM Journal on Control and Optimization*, 45(1):174–206, March 2006.
- [11] G. Chesi and Y. S. Hung. Establishing convexity of polynomial Lyapunov functions and their sublevel sets. *IEEE Transactions on Automatic Control*, 53(10):2431–2436, November 2008.
- [12] J. Desharnais, A. Edalat, and P. Panangaden. Bisimulation for labeled Markov processes. *Information and Computation*, 179(2):163–193, December 2002.
- [13] J. Desharnais, F. Laviolette, and M. Tracol. Approximate analysis of probabilistic processes: logic, simulation and games. In *Proceedings of the International Conference on Quantitative Evaluation of SysTems*, pages 264–273, September 2008.
- [14] J. J. Duistermaat and J. A. C. Kolk. *Multidimensional Real Analysis I: Differentiation*, volume 1. Cambridge University Press, 2004.
- [15] A. Girard and G. J. Pappas. Approximation metrics for discrete and continuous systems. *IEEE Transactions on Automatic Control*, 25(5):782–798, May 2007.
- [16] A. Girard, G. Pola, and P. Tabuada. Approximately bisimilar symbolic models for incrementally stable switched systems. *IEEE Transactions on Automatic Control*, 55(1):116–126, January 2010.
- [17] L. C. G. J. M. Habets, P. J. Collins, and J. H. Van Schuppen. Reachability and control synthesis for piecewise-affine hybrid systems on simplices. *IEEE Transactions on Automatic Control*, 51(6):938–948, June 2006.
- [18] G. Holzmann. *The SPIN model checker: Primer and reference manual*. Addison-Wesley Professional, 1 edition edition, 2003.
- [19] A. A. Julius and G. J. Pappas. Approximations of stochastic hybrid systems. *IEEE Transaction on Automatic Control*, 54(6):1193–1203, June 2009.
- [20] H. J. Kushner. *Stochastic stability and control*. ser. Mathematics in Science and Engineering. New York: Academic Press, 1967.
- [21] H. J. Kushner and P. G. Dupuis. *Numerical Methods for Stochastic Control Problems in Continuous Time*. Springer-Verlag, New York, 2001.
- [22] M. Lahijanian, S. B. Andersson, and C. Belta. A probabilistic approach for control of a stochastic system from LTL specifications. In *Proceedings of the 48th IEEE Conference on Decision and Control*, pages 2236–2241, December 2009.
- [23] M. Lazar and A. I. Doban. On infinity norms as lyapunov functions for continuous-time dynamical systems. In *Proceedings of the 50th IEEE Conference on Decision and Control and European Control Conference*, pages 7567–7572, December 2011.
- [24] J. Liu, U. Topcu, N. Ozay, and R. M. Murray. Reactive controllers for differentially flat systems with temporal logic constraints. In *Proceedings of the 51st IEEE Conference on Decision and Control*, pages 7664–7670, December 2012.
- [25] R. Majumdar and M. Zamani. Approximately bisimilar symbolic models for digital control systems. In M. Parthasarathy and S. A. Seshia, editors, *Computer Aided Verification (CAV)*, volume 7358 of *LNCS*, pages 362–377. Springer-Verlag, July 2012.
- [26] O. Maler, A. Pnueli, and J. Sifakis. On the synthesis of discrete controllers for timed systems. In E. W. Mayr and C. Puech, editors, *Symposium on Theoretical Aspects of Computer Science*, volume 900 of *LNCS*, pages 229–242. Springer-Verlag, 1995.
- [27] M. Mazo Jr., A. Davitian, and P. Tabuada. PESSOA: A tool for embedded control software synthesis. In T. Touili, B. Cook, and

- P. Jackson, editors, *Computer Aided Verification (CAV)*, volume 6174 of *LNCS*, pages 566–569. Springer-Verlag, July 2010.
- [28] B. Messner and D. Tilbury. Control tutorial for Matlab and Simulink. Electronically available at: <http://www.library.cmu.edu/ctms/ctms/>.
- [29] R. Milner. *Communication and Concurrency*. Prentice-Hall, Inc., 1989.
- [30] B. K. Oksendal. *Stochastic differential equations: An introduction with applications*. Springer, 5th edition, November 2002.
- [31] A. Papachristodoulou, J. Anderson, G. Valmorbida, S. Prajna, P. Seiler, and P. A. Parrilo. SOSTOOLS version 3.00 sum of squares optimization toolbox for MATLAB. *arXiv: 1310.4716*, October 2013.
- [32] A. Pavlov, N. van de Wouw, and H. Nijmeijer. *Uniform output regulation of nonlinear systems: a convergent dynamics approach*. Springer, Berlin, 2005.
- [33] G. Pola, A. Borri, and M. D. Di Benedetto. Integrated design of symbolic controllers for nonlinear systems. *IEEE Transactions on Automatic Control*, 57(2):534–539, February 2012.
- [34] G. Pola, A. Girard, and P. Tabuada. Approximately bisimilar symbolic models for nonlinear control systems. *Automatica*, 44(10):2508–2516, October 2008.
- [35] G. Pola and P. Tabuada. Symbolic models for nonlinear control systems: Alternating approximate bisimulations. *SIAM Journal on Control and Optimization*, 48(2):719–733, February 2009.
- [36] G. Reilßig. Computing abstractions of nonlinear systems. *IEEE Transactions on Automatic Control*, 56(11):2583–2598, November 2011.
- [37] J. Spiliotis and J. Tsinias. Notions of exponential robust stochastic stability, ISS and their Lyapunov characterizations. *International Journal of Robust and Nonlinear Control*, 13(12):173–187, February 2003.
- [38] J. Sproston. Discrete-time verification and control for probabilistic rectangular hybrid automata. In *Proceedings of the 8th International Conference on Quantitative Evaluation of Systems*, pages 79–88, September 2011.
- [39] P. Tabuada. *Verification and Control of Hybrid Systems, A symbolic approach*. Springer, 1st edition, June 2009.
- [40] Y. Tazaki and J. Imura. Discrete-state abstractions of nonlinear systems using multi-resolution quantizer. In *Proceedings of the 12th International Conference on Hybrid Systems: Computation and Control*, volume 5469, pages 351–365, April 2009.
- [41] J. Yeh. *Martingales and Stochastic Analysis*. Series on Multivariate Analysis. World Scientific, 1995.
- [42] M. Zamani and A. Abate. Symbolic control of stochastic switched systems via finite abstractions. In K. Joshi, M. Siegle, M. Stoelinga, and P. R. D’Argenio, editors, *Quantitative Evaluation of Systems*, volume 8054 of *Lecture Notes in Computer Science*, pages 305–321. Springer Berlin Heidelberg, August 2013.
- [43] M. Zamani and A. Abate. Approximately bisimilar symbolic models for randomly switched stochastic systems. *Systems and Control Letters, forthcoming*, 2014.
- [44] M. Zamani, P. Mohajerin Esfahani, A. Abate, and J. Lygeros. Symbolic models for stochastic control systems without stability assumptions. In *Proceedings of European Control Conference*, pages 4257–4262, July 2013.
- [45] M. Zamani, G. Pola, M. Mazo Jr., and P. Tabuada. Symbolic models for nonlinear control systems without stability assumptions. *IEEE Transactions on Automatic Control*, 57(7):1804–1809, July 2012.
- [46] M. Zamani and P. Tabuada. Backstepping design for incremental stability. *IEEE Transactions on Automatic Control*, 56(9):2184–2189, September 2011.
- [47] M. Zamani, I. Tkachev, and A. Abate. Bisimilar symbolic models for stochastic control systems without state-space discretization. In *Proceedings of the 17th International Conference on Hybrid Systems: Computation and Control*, pages 41–50. ACM New York, NY, USA, April 2014.

## IX. APPENDIX

*Proof of Theorem 3.3:* The proof is a consequence of the application of Gronwall’s inequality and of Ito’s lemma [30, pp. 80 and 123]. Assume that there exists a  $\delta$ -ISS- $M_q$  Lyapunov function in the sense of Definition 3.2. For any  $t \in \mathbb{R}_0^+$ , any  $v, v' \in \mathcal{U}$ , and any  $\mathbb{R}^n$ -valued random variables  $a$  and  $a'$  that are measurable in  $\mathcal{F}_0$ , we obtain

$$\begin{aligned} & \mathbb{E}[V(\xi_{av}(t), \xi_{a'v'}(t))] \\ &= \mathbb{E}\left[V(a, a') + \int_0^t \mathcal{L}^{v(s), v'(s)} V(\xi_{av}(s), \xi_{a'v'}(s)) ds\right] \\ &\leq \mathbb{E}\left[V(a, a') + \int_0^t (-\kappa V(\xi_{av}(s), \xi_{a'v'}(s)) + \rho(\|v(s) - v'(s)\|)) ds\right] \\ &\leq -\kappa \int_0^t \mathbb{E}[V(\xi_{av}(s), \xi_{a'v'}(s))] ds + \mathbb{E}[V(a, a')] + \rho(\|v - v'\|_\infty)t, \end{aligned}$$

which, by virtue of Gronwall’s inequality, leads to

$$\begin{aligned} \mathbb{E}[V(\xi_{av}(t), \xi_{a'v'}(t))] &\leq \mathbb{E}[V(a, a')]e^{-\kappa t} + te^{-\kappa t}\rho(\|v - v'\|_\infty) \\ &\leq \mathbb{E}[V(a, a')]e^{-\kappa t} + \frac{1}{e\kappa}\rho(\|v - v'\|_\infty). \end{aligned} \quad (\text{IX.1})$$

Hence, using property (ii) in Definition 3.2, we have

$$\begin{aligned} & \underline{\alpha}(\mathbb{E}[\|\xi_{av}(t) - \xi_{a'v'}(t)\|^q]) \leq \mathbb{E}[\underline{\alpha}(\|\xi_{av}(t) - \xi_{a'v'}(t)\|^q)] \\ &\leq \mathbb{E}[V(\xi_{av}(t), \xi_{a'v'}(t))] \leq \mathbb{E}[V(a, a')]e^{-\kappa t} + \frac{1}{e\kappa}\rho(\|v - v'\|_\infty) \\ &\leq \mathbb{E}[\bar{\alpha}(\|a - a'\|^q)]e^{-\kappa t} + \frac{1}{e\kappa}\rho(\|v - v'\|_\infty) \\ &\leq \bar{\alpha}(\mathbb{E}[\|a - a'\|^q])e^{-\kappa t} + \frac{1}{e\kappa}\rho(\|v - v'\|_\infty), \end{aligned} \quad (\text{IX.2})$$

where the first and last inequalities follow from property (i) and Jensen’s inequality [30, p. 310]. Since  $\underline{\alpha} \in \mathcal{K}_\infty$ , the inequality (IX.2) yields

$$\begin{aligned} & \mathbb{E}[\|\xi_{av}(t) - \xi_{a'v'}(t)\|^q] \\ &\leq \underline{\alpha}^{-1}\left(\bar{\alpha}(\mathbb{E}[\|a - a'\|^q])e^{-\kappa t} + \frac{1}{e\kappa}\rho(\|v - v'\|_\infty)\right) \\ &\leq \underline{\alpha}^{-1}\left(\bar{\alpha}(\mathbb{E}[\|a - a'\|^q])e^{-\kappa t} + \bar{\alpha}(\mathbb{E}[\|a - a'\|^q])e^{-\kappa t}\right) \\ &\quad + \underline{\alpha}^{-1}\left(\frac{1}{e\kappa}\rho(\|v - v'\|_\infty) + \frac{1}{e\kappa}\rho(\|v - v'\|_\infty)\right) \\ &\leq \underline{\alpha}^{-1}\left(2\bar{\alpha}(\mathbb{E}[\|a - a'\|^q])e^{-\kappa t}\right) + \underline{\alpha}^{-1}\left(\frac{2}{e\kappa}\rho(\|v - v'\|_\infty)\right). \end{aligned}$$

Therefore, by introducing functions  $\beta$  and  $\gamma$  as

$$\begin{aligned} \beta(\mathbb{E}[\|a - a'\|^q], t) &:= \underline{\alpha}^{-1}\left(2\bar{\alpha}(\mathbb{E}[\|a - a'\|^q])e^{-\kappa t}\right), \\ \gamma(\|v - v'\|_\infty) &:= \underline{\alpha}^{-1}\left(\frac{2}{e\kappa}\rho(\|v - v'\|_\infty)\right), \end{aligned}$$

condition (III.1) is satisfied. Hence, the system  $\Sigma$  is  $\delta$ -ISS- $M_q$ . Note that if  $\underline{\alpha}$  is linear, one can substitute the coefficients 2 in the aforementioned expressions of  $\beta$  and  $\gamma$  with 1 to get less conservative upper bound in (III.1). ■

*Proof of Lemma 3.4:* It is not difficult to check that the function  $V$  in (III.3) satisfies properties (i) and (ii) of Definition 3.2 with functions  $\underline{\alpha}(y) := \left(\frac{1}{q}\lambda_{\min}(P)\right)^{\frac{q}{2}}y$  and  $\bar{\alpha}(y) := \left(\frac{q}{2}\lambda_{\max}(P)\right)^{\frac{q}{2}}y$ . It then suffices to verify property (iii). We verify property (iii) for the case that  $f$  is differentiable and using condition (III.5). The proof, using condition (III.4), follows similarly by removing the inequalities in the proof including derivative of  $f$ . By the definition of  $V$  in (III.3), for any  $x, x' \in \mathbb{R}^n$  such that  $x \neq x'$ , and for  $q \in \{1, 2\}$ , one has

$$\begin{aligned} \partial_x V &= -\partial_{x'} V = (x - x')^T P \left(\tilde{V}(x, x')\right)^{\frac{q}{2}-1}, \\ \partial_{x, x} V &= \partial_{x', x'} V = -\partial_{x, x'} V \\ &= P \left(\tilde{V}(x, x')\right)^{\frac{q}{2}-1} + \frac{q-2}{q} P(x - x')(x - x')^T P \left(\tilde{V}(x, x')\right)^{\frac{q}{2}-2}. \end{aligned}$$

Therefore, following the definition of  $\mathcal{L}^{u, u'}$ , and for any  $x, x', z \in \mathbb{R}^n$  such that  $x \neq x'$ , and any  $u, u' \in \mathcal{U}$ , one obtains the chain of (in)equalities in (IX.3). In (IX.3),  $z \in \mathbb{R}^n$  and the mean value theorem [14] is applied to the differentiable function  $x \mapsto f(x, u)$  at points  $x, x'$  for a given input value  $u \in \mathcal{U}$  and  $L_u$  is the Lipschitz constant, as introduced in Definition 2.1. Therefore, the function  $V$  in (III.3) satisfies property (iii) of Definition 3.2 with positive constant  $\kappa = \frac{\tilde{\kappa}}{q}$  and  $\mathcal{K}_\infty$  function  $\rho(r) = \frac{n^{\frac{q}{2}} L_u^q}{\tilde{\kappa}^{q-1}} \left\| \sqrt{P} \right\|^q r^q$ . ■

*Proof of Lemma 3.8:* In the proof, we use the notation  $\sigma\sigma^T(x)$  instead of  $\sigma(x)\sigma^T(x)$  and  $H(V)(x, x')$  for the value of the Hessian matrix of  $V$  at  $(x, x') \in \mathbb{R}^{2n}$ . We drop the arguments of  $\partial_{x, x} V$ ,  $\partial_{x'} V$ , and  $H(V)$  for the sake of simplicity. In view of Ito’s formula, Jensen’s inequality, and similar to the calculations in Lemma 3.4, we

$$\begin{aligned}
\mathcal{L}^{u,u'} V(x, x') &= (x - x')^T P (\tilde{V}(x, x'))^{\frac{q}{2}-1} (f(x, u) - f(x', u')) + \frac{1}{2} \text{Tr} \left( \begin{bmatrix} \sigma(x) \\ \sigma(x') \end{bmatrix} \begin{bmatrix} \sigma^T(x) & \sigma^T(x') \end{bmatrix} \begin{bmatrix} \partial_{x,x} V & -\partial_{x,x} V \\ -\partial_{x,x} V & \partial_{x,x} V \end{bmatrix} \right) \\
&= (x - x')^T P (\tilde{V}(x, x'))^{\frac{q}{2}-1} (f(x, u) - f(x', u')) + \frac{1}{2} \text{Tr} \left( (\sigma(x) - \sigma(x')) (\sigma^T(x) - \sigma^T(x')) \partial_{x,x} V \right) \\
&= (x - x')^T P (\tilde{V}(x, x'))^{\frac{q}{2}-1} (f(x, u) - f(x', u')) + \frac{1}{2} \|\sqrt{P} (\sigma(x) - \sigma(x'))\|_F^2 (\tilde{V}(x, x'))^{\frac{q}{2}-2} \\
&\leq (x - x')^T P (\tilde{V}(x, x'))^{\frac{q}{2}-1} (f(x, u) - f(x', u) + f(x', u) - f(x', u')) + \frac{1}{2} \|\sqrt{P} (\sigma(x) - \sigma(x'))\|_F^2 (\tilde{V}(x, x'))^{\frac{q}{2}-1} \\
&\leq (x - x')^T P (\tilde{V}(x, x'))^{\frac{q}{2}-1} \partial_x f(z, u)(x - x') + (x - x')^T P (\tilde{V}(x, x'))^{\frac{q}{2}-1} (f(x', u) - f(x', u')) + \frac{1}{2} \|\sqrt{P} (\sigma(x) - \sigma(x'))\|_F^2 (\tilde{V}(x, x'))^{\frac{q}{2}-1} \\
&\leq \left( (x - x')^T P \partial_x f(z, u)(x - x') + \frac{1}{2} \|\sqrt{P} (\sigma(x) - \sigma(x'))\|_F^2 \right) (\tilde{V}(x, x'))^{\frac{q}{2}-1} + \sqrt{n} \left( (x - x')^T P (x - x') \right)^{\frac{1}{2}} \|\sqrt{P}\|_{L_u} \|u - u'\| (\tilde{V}(x, x'))^{\frac{q}{2}-1} \\
&\leq -\tilde{\kappa} V(x, x') + (q-1) \frac{\tilde{\kappa}}{2} V(x, x') + \frac{\frac{q}{2} L_u^q}{\tilde{\kappa}^{q-1}} \|\sqrt{P}\|^q \|u - u'\|^q = -\frac{\tilde{\kappa}}{q} V(x, x') + \frac{\frac{q}{2} L_u^q}{\tilde{\kappa}^{q-1}} \|\sqrt{P}\|^q \|u - u'\|^q. \tag{IX.3}
\end{aligned}$$

have

$$\begin{aligned}
\underline{\alpha} \left( \mathbb{E} \left[ \left\| \xi_{xv}(t) - \bar{\xi}_{xv}(t) \right\|^q \right] \right) &\leq \mathbb{E} \left[ \underline{\alpha} \left( \left\| \xi_{xv}(t) - \bar{\xi}_{xv}(t) \right\|^q \right) \right] \\
&\leq \mathbb{E} \left[ V \left( \xi_{xv}(t), \bar{\xi}_{xv}(t) \right) \right] = \int_0^t \mathbb{E} \left[ \mathcal{L}^{v(s), v(s)} V \left( \xi_{xv}(s), \bar{\xi}_{xv}(s) \right) \right] ds \\
&= \int_0^t \mathbb{E} \left[ \begin{bmatrix} \partial_x V & \partial_{x'} V \end{bmatrix} \begin{bmatrix} f(\xi_{xv}(s), v(s)) \\ f(\bar{\xi}_{xv}(s), v(s)) \end{bmatrix} \right. \\
&\quad \left. + \frac{1}{2} \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) \partial_{x,x} V \right) \right] ds \\
&= \int_0^t \mathbb{E} \left[ \begin{bmatrix} \partial_x V & \partial_{x'} V \end{bmatrix} \begin{bmatrix} f(\xi_{xv}(s), v(s)) \\ f(\bar{\xi}_{xv}(s), v(s)) \end{bmatrix} \right. \\
&\quad \left. + \frac{1}{2} \text{Tr} \left( \begin{bmatrix} \sigma(\xi_{xv}(s)) \\ \sigma(\bar{\xi}_{xv}(s)) \end{bmatrix} \begin{bmatrix} \sigma^T(\xi_{xv}(s)) & \sigma^T(\bar{\xi}_{xv}(s)) \end{bmatrix} H(V) \right) \right. \\
&\quad \left. + \frac{1}{2} \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) \partial_{x,x} V \right) \right. \\
&\quad \left. - \frac{1}{2} \text{Tr} \left( \begin{bmatrix} \sigma(\xi_{xv}(s)) \\ \sigma(\bar{\xi}_{xv}(s)) \end{bmatrix} \begin{bmatrix} \sigma^T(\xi_{xv}(s)) & \sigma^T(\bar{\xi}_{xv}(s)) \end{bmatrix} H(V) \right) \right] ds \tag{IX.4}
\end{aligned}$$

$$\begin{aligned}
&\leq \int_0^t -\tilde{\kappa} \mathbb{E} \left[ V \left( \xi_{xv}(s), \bar{\xi}_{xv}(s) \right) \right] ds \tag{IX.5} \\
&\quad + \int_0^t \mathbb{E} \left[ \frac{1}{2} \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) \partial_{x,x} V \right) \right. \\
&\quad \left. - \frac{1}{2} \text{Tr} \left( \begin{bmatrix} \sigma(\xi_{xv}(s)) \\ \sigma(\bar{\xi}_{xv}(s)) \end{bmatrix} \begin{bmatrix} \sigma^T(\xi_{xv}(s)) & \sigma^T(\bar{\xi}_{xv}(s)) \end{bmatrix} H(V) \right) \right] ds \\
&\leq \int_0^t -\tilde{\kappa} \mathbb{E} \left[ V \left( \xi_{xv}(s), \bar{\xi}_{xv}(s) \right) \right] ds \tag{IX.6} \\
&\quad + \int_0^t \mathbb{E} \left[ \frac{1}{2} \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) \partial_{x,x} V \right) \right] ds \leq \hat{h}(\sigma, t) e^{-\tilde{\kappa} t},
\end{aligned}$$

where the function  $\hat{h}$  can be computed as  $\hat{h}(\sigma, t) := \int_0^t \mathbb{E} \left[ \frac{1}{2} \|\sqrt{\partial_{x,x} V} \sigma(\xi_{xv}(s))\|_F^2 \right] ds$ . Inequality (IX.5) is a straightforward consequence of  $V$  satisfying the condition (iii) in Definition 3.2, and (IX.6) follows from Gronwall's inequality. Using the Lipschitz continuity assumption on the diffusion term  $\sigma$ , we get:

$$\begin{aligned}
&\int_0^t \mathbb{E} \left[ \frac{1}{2} \|\sqrt{\partial_{x,x} V} \sigma(\xi_{xv}(s))\|_F^2 \right] ds \leq \\
&\frac{1}{2} \|\sqrt{P}\|^2 n \min\{n, p\} Z^2 \int_0^t \mathbb{E} \left[ \|\xi_{xv}(s)\|^2 \right] ds.
\end{aligned}$$

Since  $V$  is a  $\delta$ -ISS- $M_q$  Lyapunov function,  $q \geq 2$ ,  $f(0_n, 0_m) = 0_n$ ,  $\sigma(0_n) = 0_{n \times p}$ , and using functions  $\beta$  and  $\gamma$  in (III.1), one can verify that:

$$\begin{aligned}
&\left( \mathbb{E} \left[ \|\xi_{xv}(t)\|^2 \right] \right)^{\frac{q}{2}} \leq \mathbb{E} \left[ \left( \|\xi_{xv}(t)\|^2 \right)^{\frac{q}{2}} \right] = \mathbb{E} \left[ \|\xi_{xv}(t)\|^q \right] \\
&\leq \beta (\|x\|^q, t) + \gamma (\|u\|_\infty) \leq \beta \left( \sup_{x \in \mathbb{D}} \{\|x\|^q\}, t \right) + \gamma \left( \sup_{u \in \mathbb{U}} \{\|u\|\} \right),
\end{aligned}$$

hence,

$$\mathbb{E} \left[ \|\xi_{xv}(t)\|^2 \right] \leq \left( \beta \left( \sup_{x \in \mathbb{D}} \{\|x\|^q\}, t \right) + \gamma \left( \sup_{u \in \mathbb{U}} \{\|u\|\} \right) \right)^{\frac{2}{q}}.$$

Therefore, we have

$$\begin{aligned}
\hat{h}(\sigma, t) &\leq \frac{1}{2} \|\sqrt{P}\|^2 n \min\{n, p\} Z^2 \\
&\quad \cdot \int_0^t \left( \beta \left( \sup_{x \in \mathbb{D}} \{\|x\|^q\}, s \right) + \gamma \left( \sup_{u \in \mathbb{U}} \{\|u\|\} \right) \right)^{\frac{2}{q}} ds.
\end{aligned}$$

By defining:

$$\begin{aligned}
h(\sigma, t) &= \underline{\alpha}^{-1} \left( \frac{1}{2} \|\sqrt{P}\|^2 n \min\{n, p\} Z^2 e^{-\tilde{\kappa} t} \right. \\
&\quad \left. \cdot \int_0^t \left( \beta \left( \sup_{x \in \mathbb{D}} \{\|x\|^q\}, s \right) + \gamma \left( \sup_{u \in \mathbb{U}} \{\|u\|\} \right) \right)^{\frac{2}{q}} ds \right), \tag{IX.7}
\end{aligned}$$

we obtain  $\mathbb{E} \left[ \|\xi_{xv}(t) - \bar{\xi}_{xv}(t)\|^q \right] \leq h(\sigma, t)$ . It is not hard to observe that the proposed function  $h$  meets the conditions of the lemma. ■

*Proof of Lemma 3.10:* In the proof, we use the notation  $\sigma \sigma^T(x)$  instead of  $\sigma(x) \sigma^T(x)$  for the sake of simplicity. In view of Ito's formula and similar to the calculations in Lemma 3.8, we have

$$\begin{aligned}
&\frac{1}{2} \lambda_{\min}(P) \mathbb{E} \left[ \|\xi_{xv}(t) - \bar{\xi}_{xv}(t)\|^2 \right] \leq \frac{q}{2} \mathbb{E} \left[ V^{\frac{2}{q}} \left( \xi_{xv}(t), \bar{\xi}_{xv}(t) \right) \right] \\
&= \int_0^t \mathbb{E} \left[ \left( \xi_{xv}(s) - \bar{\xi}_{xv}(s) \right)^T P \left( f(\xi_{xv}(s), v(s)) - f(\bar{\xi}_{xv}(s), v(s)) \right) \right. \\
&\quad \left. + \frac{1}{2} \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) P \right) \right] ds \\
&= \int_0^t \mathbb{E} \left[ \left( \xi_{xv}(s) - \bar{\xi}_{xv}(s) \right)^T P \left( f(\xi_{xv}(s), v(s)) - f(\bar{\xi}_{xv}(s), v(s)) \right) \right. \\
&\quad \left. + \frac{1}{2} \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s) - \bar{\xi}_{xv}(s)) P \right) \right. \\
&\quad \left. + \frac{1}{2} \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) P - \sigma \sigma^T (\xi_{xv}(s) - \bar{\xi}_{xv}(s)) P \right) \right] ds \\
&\leq \int_0^t -\tilde{\kappa} \mathbb{E} \left[ V^{\frac{2}{q}} \left( \xi_{xv}(s), \bar{\xi}_{xv}(s) \right) \right] ds \tag{IX.8} \\
&\quad + \frac{1}{2} \int_0^t \mathbb{E} \left[ \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) P - \sigma \sigma^T (\xi_{xv}(s) - \bar{\xi}_{xv}(s)) P \right) \right] ds \\
&\leq \int_0^t -\tilde{\kappa} \mathbb{E} \left[ V^{\frac{2}{q}} \left( \xi_{xv}(s), \bar{\xi}_{xv}(s) \right) \right] ds \tag{IX.9} \\
&\quad + \frac{1}{2} \int_0^t \mathbb{E} \left[ \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) P \right) \right] ds \leq \hat{h}(\sigma, t) e^{-\frac{2\tilde{\kappa} t}{q}},
\end{aligned}$$

where the function  $\hat{h}$  can be computed as  $\hat{h}(\sigma, t) := \frac{1}{2} \int_0^t \mathbb{E} \left[ \|\sqrt{P} \sigma(\xi_{xv}(s))\|_F^2 \right] ds$ . Inequality (IX.8) is a straightforward consequence of (III.4), and (IX.9) follows from Gronwall's inequality. Using the Lipschitz continuity assumption on the diffusion term  $\sigma$ ,

we get:

$$\begin{aligned} & \frac{1}{2} \int_0^t \mathbb{E} \left[ \left\| \sqrt{P} \sigma(\xi_{xv}(s)) \right\|_F^2 \right] ds \\ & \leq \frac{1}{2} \left\| \sqrt{P} \right\|^2 n \min\{n, p\} Z^2 \int_0^t \mathbb{E} \left[ \|\xi_{xv}(s)\|^2 \right] ds. \end{aligned}$$

Since  $V^{\frac{2}{q}}$  is a  $\delta$ -ISS- $M_2$  Lyapunov function,  $f(0_n, 0_m) = 0_n$ ,  $\sigma(0_n) = 0_{n \times p}$ , and using functions  $\beta$  and  $\gamma$  in (III.1), obtained by the  $\delta$ -ISS- $M_q$  Lyapunov function  $V^{\frac{2}{q}}$ , one can verify that:

$$\mathbb{E} \left[ \|\xi_{xv}(t)\|^2 \right] \leq \frac{n \lambda_{\max}(P)}{\lambda_{\min}(P)} \|x\|^2 e^{-\frac{\tilde{\kappa}t}{2}} + \frac{2n \left\| \sqrt{P} \right\|^2 L_u^2}{e \tilde{\kappa}^2 \lambda_{\min}(P)} \|v\|^2.$$

Therefore, one obtains:

$$\begin{aligned} \hat{h}(\sigma, t) & \leq \frac{\left\| \sqrt{P} \right\|^2 n^2 \min\{n, p\} Z^2}{\lambda_{\min}(P) \tilde{\kappa}} \\ & \left( \lambda_{\max}(P) \left( 1 - e^{-\frac{\tilde{\kappa}t}{2}} \right) \sup_{x \in \mathbb{D}} \{\|x\|^2\} + \frac{\left\| \sqrt{P} \right\|^2 L_u^2}{e \tilde{\kappa}} \sup_{u \in \mathbb{U}} \{\|u\|^2\} t \right). \end{aligned}$$

By defining:

$$\begin{aligned} h(\sigma, t) & = \frac{2 \left\| \sqrt{P} \right\|^2 n^2 \min\{n, p\} Z^2 e^{-\frac{2\tilde{\kappa}t}{q}}}{\lambda_{\min}^2(P) \tilde{\kappa}} \\ & \left( \lambda_{\max}(P) \left( 1 - e^{-\frac{\tilde{\kappa}t}{2}} \right) \sup_{x \in \mathbb{D}} \{\|x\|^2\} + \frac{\left\| \sqrt{P} \right\|^2 L_u^2}{e \tilde{\kappa}} \sup_{u \in \mathbb{U}} \{\|u\|^2\} t \right), \end{aligned} \quad (\text{IX.10})$$

we obtain  $\mathbb{E} \left[ \|\xi_{xv}(t) - \bar{\xi}_{xv}(t)\|^2 \right] \leq h(\sigma, t)$ . It is not hard to observe that the proposed function  $h$  meets the conditions of the lemma. ■

*Proof of Corollary 3.11:* Motivated by inequality (IX.8), one can obtain

$$\begin{aligned} & \mathbb{E} \left[ \text{Tr} \left( \sigma \sigma^T (\xi_{xv}(s)) P - \sigma \sigma^T (\xi_{xv}(s) - \bar{\xi}_{xv}(s)) P \right) \right] \\ & = \mathbb{E} \left[ \xi_{xv}^T(s) \left( \sum_{i=1}^p \sigma_i^T P \sigma_i \right) \xi_{xv}(s) \right. \\ & \quad \left. - (\xi_{xv}(s) - \bar{\xi}_{xv}(s))^T \left( \sum_{i=1}^p \sigma_i^T P \sigma_i \right) (\xi_{xv}(s) - \bar{\xi}_{xv}(s)) \right] \\ & = \bar{\xi}_{xv}^T(s) \left( \sum_{i=1}^p \sigma_i^T P \sigma_i \right) \bar{\xi}_{xv}(s) \leq n \lambda_{\max} \left( \sum_{i=1}^p \sigma_i^T P \sigma_i \right) \|\bar{\xi}_{xv}(s)\|^2. \end{aligned}$$

It can be readily verified that

$$\begin{aligned} \|\bar{\xi}_{xv}(t)\| & \leq \|e^{At}\| \|x\| + \left( \int_0^t \|e^{As} B\| ds \right) \|v\|_{\infty} \\ & \leq \|e^{At}\| \sup_{x \in \mathbb{D}} \{\|x\|\} + \left( \int_0^t \|e^{As} B\| ds \right) \sup_{u \in \mathbb{U}} \{\|u\|\}. \end{aligned} \quad (\text{IX.11})$$

The above approximation, together with (IX.8), leads to a more explicit bound in terms of the system parameters as follows:

$$\begin{aligned} \mathbb{E} \left[ \|\xi_{xv}(t) - \bar{\xi}_{xv}(t)\|^2 \right] & \leq \frac{n \lambda_{\max} \left( \sum_{i=1}^p \sigma_i^T P \sigma_i \right) e^{-\tilde{\kappa}t}}{\lambda_{\min}(P)} \\ & \int_0^t \left( \|e^{As}\| \sup_{x \in \mathbb{D}} \{\|x\|\} + \left( \int_0^s \|e^{Ar} B\| dr \right) \sup_{u \in \mathbb{U}} \{\|u\|\} \right)^2 ds, \end{aligned} \quad (\text{IX.12})$$

where  $\hat{\kappa} = \frac{2\tilde{\kappa}}{q}$ . ■

*Proof of Theorem 5.1:* We start by proving that  $S_{\tau}(\Sigma) \preceq_{\mathcal{S}}^{\varepsilon} S_q(\Sigma)$ . Consider the relation  $R \subseteq X_{\tau} \times X_q$  defined by  $(x_{\tau}, x_q) \in R$  if and only if  $\mathbb{E}[V(\bar{H}_{\tau}(x_{\tau}), H_q(x_q))] = \mathbb{E}[V(x'_{\tau}, x'_q)] \leq \underline{\alpha}(\varepsilon^q)$ . Since  $X_{\tau_0} \subseteq \bigcup_{p \in [\mathbb{R}^n]_{\eta}} \mathcal{B}_{\eta}(p)$ , for every  $x_{\tau_0} \in X_{\tau_0}$  there always exists  $x_{q_0} \in X_{q_0}$  such that  $\|x_{\tau_0} - x_{q_0}\| \leq \eta$ .

Then,

$$\begin{aligned} \mathbb{E}[V(x_{\tau_0}, x_{q_0})] & = V(x_{\tau_0}, x_{q_0}) \leq \bar{\alpha}(\|x_{\tau_0} - x_{q_0}\|^q) \\ & \leq \bar{\alpha}(\eta^q) \leq \underline{\alpha}(\varepsilon^q), \end{aligned}$$

because of (V.3) and since  $\bar{\alpha}$  is a  $\mathcal{K}_{\infty}$  function. Hence,  $(x_{\tau_0}, x_{q_0}) \in R$  and condition (i) in Definition 4.2 is satisfied. Now consider any  $(x_{\tau}, x_q) \in R$ . Condition (ii) in Definition 4.2 is satisfied because

$$\left( \mathbb{E}[\|x_{\tau} - x_q\|^q] \right)^{\frac{1}{q}} \leq \left( \underline{\alpha}^{-1}(\mathbb{E}[V(x_{\tau}, x_q)]) \right)^{\frac{1}{q}} \leq \varepsilon. \quad (\text{IX.13})$$

We used the convexity assumption of  $\underline{\alpha}$  and the Jensen inequality [30] to show the inequalities in (IX.13). Let us now show that condition (iii) in Definition 4.2 holds. Consider any  $v_{\tau} \in U_{\tau}$ . Choose an input  $u_q \in U_q$  satisfying

$$\|v_{\tau} - u_q\|_{\infty} = \|v_{\tau}(0) - u_q(0)\| \leq \mu. \quad (\text{IX.14})$$

Note that the existence of such  $u_q$  is guaranteed by  $\mathbb{U}$  being a finite union of boxes and by the inequality  $\mu \leq \text{span}(\mathbb{U})$  which guarantees that  $\mathbb{U} \subseteq \bigcup_{p \in [\mathbb{U}]_{\mu}} \mathcal{B}_{\mu}(p)$ . Consider the transition  $x_{\tau} \xrightarrow{v_{\tau}} x'_{\tau} = \xi_{x_{\tau} v_{\tau}}(\tau)$  P-a.s. in  $S_{\tau}(\Sigma)$ . Since  $V$  is a  $\delta$ -ISS- $M_q$  Lyapunov function for  $\Sigma$ , in light of (IX.1) as well as (IX.14), we have

$$\begin{aligned} \mathbb{E}[V(x'_{\tau}, \xi_{x_q u_q}(\tau))] & \leq \mathbb{E}[V(x_{\tau}, x_q)] e^{-\kappa\tau} + \frac{1}{e\kappa} \rho(\|v_{\tau} - u_q\|_{\infty}) \\ & \leq \underline{\alpha}(\varepsilon^q) e^{-\kappa\tau} + \frac{1}{e\kappa} \rho(\mu). \end{aligned} \quad (\text{IX.15})$$

Since  $\mathbb{R}^n \subseteq \bigcup_{p \in [\mathbb{R}^n]_{\eta}} \mathcal{B}_{\eta}(p)$ , there exists a  $x'_q \in X_q$  such that

$$\|\bar{\xi}_{x_q u_q}(\tau) - x'_q\| \leq \eta, \quad (\text{IX.16})$$

which, by the definition of  $S_q(\Sigma)$ , implies the existence of  $x_q \xrightarrow{u_q} x'_q$  in  $S_q(\Sigma)$ . Using Lemmas 3.8 or 3.10, the concavity of  $\hat{\gamma}$ , the Jensen inequality [30], the inequalities (V.2), (V.4), (IX.15), (IX.16), and triangle inequality, we obtain

$$\begin{aligned} & \mathbb{E}[V(x'_{\tau}, x'_q)] \\ & = \mathbb{E}[V(x'_{\tau}, \xi_{x_q u_q}(\tau)) + V(x'_{\tau}, x'_q) - V(x'_{\tau}, \xi_{x_q u_q}(\tau))] \\ & = \mathbb{E}[V(x'_{\tau}, \xi_{x_q u_q}(\tau))] + \mathbb{E}[V(x'_{\tau}, x'_q) - V(x'_{\tau}, \xi_{x_q u_q}(\tau))] \\ & \leq \underline{\alpha}(\varepsilon^q) e^{-\kappa\tau} + \frac{1}{e\kappa} \rho(\mu) + \mathbb{E}[\hat{\gamma}(\|\xi_{x_q u_q}(\tau) - x'_q\|)] \\ & \leq \underline{\alpha}(\varepsilon^q) e^{-\kappa\tau} + \frac{1}{e\kappa} \rho(\mu) \\ & \quad + \hat{\gamma} \left( \mathbb{E} \left[ \|\xi_{x_q u_q}(\tau) - \bar{\xi}_{x_q u_q}(\tau) + \bar{\xi}_{x_q u_q}(\tau) - x'_q\| \right] \right) \\ & \leq \underline{\alpha}(\varepsilon^q) e^{-\kappa\tau} + \frac{1}{e\kappa} \rho(\mu) \\ & \quad + \hat{\gamma} \left( \mathbb{E} \left[ \|\xi_{x_q u_q}(\tau) - \bar{\xi}_{x_q u_q}(\tau)\| \right] + \|\bar{\xi}_{x_q u_q}(\tau) - x'_q\| \right) \\ & \leq \underline{\alpha}(\varepsilon^q) e^{-\kappa\tau} + \frac{1}{e\kappa} \rho(\mu) + \hat{\gamma} \left( (h(\sigma, \tau))^{\frac{1}{q}} + \eta \right) \leq \underline{\alpha}(\varepsilon^q). \end{aligned}$$

Therefore, we conclude that  $(x'_{\tau}, x'_q) \in R$  and that condition (iii) in Definition 4.2 holds.

Now we prove  $S_q(\Sigma) \preceq_{\mathcal{S}}^{\varepsilon} S_{\tau}(\Sigma)$  implying that  $R^{-1}$  is a suitable  $\varepsilon$ -approximate simulation relation from  $S_q(\Sigma)$  to  $S_{\tau}(\Sigma)$ . Consider the relation  $R \subseteq X_{\tau} \times X_q$ , defined in the first part of the proof. For every  $x_{q_0} \in X_{q_0}$ , by choosing  $x_{\tau_0} = x_{q_0}$ , we have  $V(x_{\tau_0}, x_{q_0}) = 0$  and  $(x_{\tau_0}, x_{q_0}) \in R$  and condition (i) in Definition 4.2 is satisfied. Now consider any  $(x_{\tau}, x_q) \in R$ . Condition (ii) in Definition 4.2 is satisfied because

$$\left( \mathbb{E}[\|x_{\tau} - x_q\|^q] \right)^{\frac{1}{q}} \leq \left( \underline{\alpha}^{-1}(\mathbb{E}[V(x_{\tau}, x_q)]) \right)^{\frac{1}{q}} \leq \varepsilon. \quad (\text{IX.17})$$

We used the convexity assumption of  $\underline{\alpha}$  and the Jensen inequality [30] to show the inequalities in (IX.17). Let us now show that condition (iii) in Definition 4.2 holds. Consider any  $u_q \in U_q$ . Choose the input  $v_{\tau} = u_q$  and consider  $x'_{\tau} = \xi_{x_{\tau} v_{\tau}}(\tau)$  P-a.s. in  $S_{\tau}(\Sigma)$ . Since  $V$  is a  $\delta$ -ISS- $M_q$  Lyapunov function for  $\Sigma$ , one obtains:

$$\mathbb{E}[V(x'_{\tau}, \xi_{x_q u_q}(\tau))] \leq e^{-\kappa\tau} \mathbb{E}[V(x_{\tau}, x_q)] \leq e^{-\kappa\tau} \underline{\alpha}(\varepsilon^q). \quad (\text{IX.18})$$

Using Lemmas 3.8 or 3.10, the definition of  $S_q(\Sigma)$ , the concavity of  $\hat{\gamma}$ , the Jensen inequality [30], the inequalities (V.2), (V.4), (IX.18), and the triangle inequality, we obtain

$$\begin{aligned}
& \mathbb{E} [V(x'_\tau, x'_q)] \\
&= \mathbb{E} [V(x'_\tau, \xi_{x_q u_q}(\tau)) + V(x'_\tau, x'_q) - V(x'_\tau, \xi_{x_q u_q}(\tau))] \\
&= \mathbb{E} [V(x'_\tau, \xi_{x_q u_q}(\tau))] + \mathbb{E} [V(x'_\tau, x'_q) - V(x'_\tau, \xi_{x_q u_q}(\tau))] \\
&\leq e^{-\kappa\tau} \underline{\alpha}(\varepsilon^q) + \mathbb{E} [\hat{\gamma}(\|\xi_{x_q u_q}(\tau) - x'_q\|)] \\
&\leq e^{-\kappa\tau} \underline{\alpha}(\varepsilon^q) + \hat{\gamma} \left( \mathbb{E} \left[ \left\| \xi_{x_q u_q}(\tau) - \bar{\xi}_{x_q u_q}(\tau) + \bar{\xi}_{x_q u_q}(\tau) - x'_q \right\|^q \right] \right)^{\frac{1}{q}} \\
&\leq e^{-\kappa\tau} \underline{\alpha}(\varepsilon^q) + \hat{\gamma} \left( \mathbb{E} \left[ \left\| \xi_{x_q u_q}(\tau) - \bar{\xi}_{x_q u_q}(\tau) \right\|^q \right] + \left\| \bar{\xi}_{x_q u_q}(\tau) - x'_q \right\|^q \right)^{\frac{1}{q}} \\
&\leq e^{-\kappa\tau} \underline{\alpha}(\varepsilon^q) + \hat{\gamma} \left( (h(\sigma, \tau))^{\frac{1}{q}} + \eta \right) \leq \underline{\alpha}(\varepsilon^q).
\end{aligned}$$

Therefore, we conclude that  $(x'_\tau, x'_q) \in R$  and condition (iii) in Definition 4.2 holds. ■

*Proof of Theorem 5.3:* We start by proving  $S_\tau(\Sigma) \preceq_S^\varepsilon S_q(\Sigma)$ . Consider the relation  $R \subseteq X_\tau \times X_q$  defined by  $(x_\tau, x_q) \in R$  if and only if  $(\mathbb{E}[\|H_\tau(x_\tau) - H_q(x_q)\|^q])^{\frac{1}{q}} = (\mathbb{E}[\|x_\tau - x_q\|^q])^{\frac{1}{q}} \leq \varepsilon$ . Since  $X_{\tau 0} \subseteq \bigcup_{p \in [\mathbb{R}^n]_\eta} \mathcal{B}_\eta(p)$ , for every  $x_{\tau 0} \in X_{\tau 0}$  there always exists  $x_{q 0} \in X_{q 0}$  such that  $\|x_{\tau 0} - x_{q 0}\| \leq \eta$ . Then,

$$(\mathbb{E}[\|x_{\tau 0} - x_{q 0}\|^q])^{\frac{1}{q}} = (\|x_{\tau 0} - x_{q 0}\|)^{\frac{1}{q}} \leq \eta \leq \varepsilon,$$

because of (V.8). Hence,  $(x_{\tau 0}, x_{q 0}) \in R$  and condition (i) in Definition 4.2 is satisfied. Now consider any  $(x_\tau, x_q) \in R$ . Condition (ii) in Definition 4.2 is satisfied by the definition of  $R$ . Let us now show that condition (iii) in Definition 4.2 holds. Consider any  $v_\tau \in U_\tau$ . Choose an input  $u_q \in U_q$  satisfying

$$\|v_\tau - u_q\|_\infty = \|v_\tau(0) - u_q(0)\| \leq \mu. \quad (\text{IX.19})$$

Note that the existence of such  $u_q$  is guaranteed by  $\mathbf{U}$  being a finite union of boxes and by the inequality  $\mu \leq \text{span}(\mathbf{U})$  which guarantees that  $\mathbf{U} \subseteq \bigcup_{p \in [\mathbf{U}]_\mu} \mathcal{B}_\mu(p)$ . Consider the transition

$x_\tau \xrightarrow{v_\tau} x'_\tau = \xi_{x_\tau v_\tau}(\tau)$  P-a.s. in  $S_\tau(\Sigma)$ . It follows from the  $\delta$ -ISS- $M_q$  assumption on  $\Sigma$  and (IX.19) that:

$$\begin{aligned}
\mathbb{E} [\|x'_\tau - \xi_{x_q u_q}(\tau)\|^q] &\leq \beta (\mathbb{E}[\|x_\tau - x_q\|^q], \tau) + \gamma(\|v_\tau - u_q\|_\infty) \\
&\leq \beta(\varepsilon^q, \tau) + \gamma(\mu).
\end{aligned} \quad (\text{IX.20})$$

Since  $\mathbb{R}^n \subseteq \bigcup_{p \in [\mathbb{R}^n]_\eta} \mathcal{B}_\eta(p)$ , there exists  $x'_q \in X_q$  such that

$$\left\| \bar{\xi}_{x_q u_q}(\tau) - x'_q \right\| \leq \eta, \quad (\text{IX.21})$$

which, by the definition of  $S_q(\Sigma)$ , implies the existence of  $x_q \xrightarrow{u_q} x'_q$  in  $S_q(\Sigma)$ . Using Lemmas 3.8 or 3.10, (V.8), (IX.20), (IX.21), and triangle inequality, we obtain

$$\begin{aligned}
& (\mathbb{E}[\|x'_\tau - x'_q\|^q])^{\frac{1}{q}} \\
&= \left( \mathbb{E} \left[ \left\| x'_\tau - \xi_{x_q u_q}(\tau) + \xi_{x_q u_q}(\tau) - \bar{\xi}_{x_q u_q}(\tau) + \bar{\xi}_{x_q u_q}(\tau) - x'_q \right\|^q \right] \right)^{\frac{1}{q}} \\
&\leq (\mathbb{E}[\|x'_\tau - \xi_{x_q u_q}(\tau)\|^q])^{\frac{1}{q}} + \left( \mathbb{E} \left[ \left\| \xi_{x_q u_q}(\tau) - \bar{\xi}_{x_q u_q}(\tau) \right\|^q \right] \right)^{\frac{1}{q}} \\
&\quad + \left( \mathbb{E} \left[ \left\| \bar{\xi}_{x_q u_q}(\tau) - x'_q \right\|^q \right] \right)^{\frac{1}{q}} \\
&\leq (\beta(\varepsilon^q, \tau) + \gamma(\mu))^{\frac{1}{q}} + (h(\sigma, \tau))^{\frac{1}{q}} + \eta \leq \varepsilon.
\end{aligned}$$

Therefore, we conclude that  $(x'_\tau, x'_q) \in R$  and that condition (iii) in Definition 4.2 holds.

Now we prove  $S_q(\Sigma) \preceq_S^\varepsilon S_\tau(\Sigma)$  implying that  $R^{-1}$  is a suitable  $\varepsilon$ -approximate simulation relation from  $S_q(\Sigma)$  to  $S_\tau(\Sigma)$ . Consider the relation  $R \subseteq X_\tau \times X_q$ , defined in the first part of the proof. For every  $x_{q 0} \in X_{q 0}$ , by choosing  $x_{\tau 0} = x_{q 0}$ , we have  $\|x_{\tau 0} - x_{q 0}\|^q = 0$  and  $(x_{\tau 0}, x_{q 0}) \in R$  and condition (i) in Definition 4.2 is satisfied. Now consider any  $(x_\tau, x_q) \in R$ . Condition (ii) in Definition 4.2 is satisfied by the definition of  $R$ . Let us now show that condition (iii) in Definition 4.2 holds. Consider any  $u_q \in U_q$ . Choose the input  $v_\tau = u_q$  and consider  $x'_\tau = \xi_{x_\tau v_\tau}(\tau)$  P-a.s. in  $S_\tau(\Sigma)$ . Since  $\Sigma$  is  $\delta$ -ISS- $M_q$ , one obtains:

$$\mathbb{E} [\|x'_\tau - \xi_{x_q u_q}(\tau)\|^q] \leq \beta (\mathbb{E}[\|x_\tau - x_q\|^q], \tau) \leq \beta(\varepsilon^q, \tau). \quad (\text{IX.22})$$

Using Lemmas 3.8 or 3.10, the definition of  $S_q(\Sigma)$ , (V.8), (IX.22), and the triangle inequality, we obtain

$$\begin{aligned}
& (\mathbb{E}[\|x'_\tau - x'_q\|^q])^{\frac{1}{q}} \\
&= \left( \mathbb{E} \left[ \left\| x'_\tau - \xi_{x_q u_q}(\tau) + \xi_{x_q u_q}(\tau) - \bar{\xi}_{x_q u_q}(\tau) + \bar{\xi}_{x_q u_q}(\tau) - x'_q \right\|^q \right] \right)^{\frac{1}{q}} \\
&\leq (\mathbb{E}[\|x'_\tau - \xi_{x_q u_q}(\tau)\|^q])^{\frac{1}{q}} + \left( \mathbb{E} \left[ \left\| \xi_{x_q u_q}(\tau) - \bar{\xi}_{x_q u_q}(\tau) \right\|^q \right] \right)^{\frac{1}{q}} \\
&\quad + \left( \mathbb{E} \left[ \left\| \bar{\xi}_{x_q u_q}(\tau) - x'_q \right\|^q \right] \right)^{\frac{1}{q}} \leq (\beta(\varepsilon^q, \tau))^{\frac{1}{q}} + (h(\sigma, \tau))^{\frac{1}{q}} + \eta \leq \varepsilon.
\end{aligned}$$

Therefore, we conclude that  $(x'_\tau, x'_q) \in R$  and condition (iii) in Definition 4.2 holds. ■

*Proof of Proposition 5.7:* Following the definition of stochastic bisimulation function in [19], for any input signal  $v \in \mathcal{U}_\tau$ , there exists an input signal  $\bar{v} \in \mathcal{U}_\tau$  such that  $\phi(\xi_{x_{\tau 0} v}, \bar{\xi}_{x_{\tau 0} \bar{v}})$  is a nonnegative supermartingale [30, Appendix C]. Furthermore, using the chosen triple  $\mathbf{q} = (\tau, \eta, \mu)$  of quantization parameters, we have:  $S_\tau(\bar{\Sigma}) \cong_S^\varepsilon S_q(\Sigma)$ , implying that for a state run  $\bar{\xi}_{x_{\tau 0} \bar{v}}$  of  $S_\tau(\bar{\Sigma})$ , there exists a state run  $X_{x_{q 0} u}$  of  $S_q(\Sigma)$  such that:

$$\sup_{k \in \mathbb{N}_0} \left\| \bar{\xi}_{x_{\tau 0} \bar{v}}(k\tau) - X_{x_{q 0} u}(k) \right\| \leq \varepsilon.$$

The above statement follows by direct application of Definition 4.2. Since an increasing concave function of a supermartingale is a supermartingale [41, Theorem 5.11], we have the following chain of (in)equalities:

$$\begin{aligned}
& \mathbb{P} \left\{ \sup_{k \in \mathbb{N}_0} \left\| \xi_{x_{\tau 0} v}(k\tau) - X_{x_{q 0} u}(k) \right\| > \varepsilon \mid x_{\tau 0} \right\} = \\
& \mathbb{P} \left\{ \sup_{k \in \mathbb{N}_0} \left\| \xi_{x_{\tau 0} v}(k\tau) - \bar{\xi}_{x_{\tau 0} \bar{v}}(k\tau) \right. \right. \\
& \quad \left. \left. + \bar{\xi}_{x_{\tau 0} \bar{v}}(k\tau) - X_{x_{q 0} u}(k) \right\| > \varepsilon \mid x_{\tau 0} \right\} \leq \\
& \mathbb{P} \left\{ \sup_{k \in \mathbb{N}_0} \left\{ \left\| \xi_{x_{\tau 0} v}(k\tau) - \bar{\xi}_{x_{\tau 0} \bar{v}}(k\tau) \right\| \right. \right. \\
& \quad \left. \left. + \left\| \bar{\xi}_{x_{\tau 0} \bar{v}}(k\tau) - X_{x_{q 0} u}(k) \right\| \right\} > \varepsilon \mid x_{\tau 0} \right\} \leq \\
& \mathbb{P} \left\{ \sup_{k \in \mathbb{N}_0} \left\{ \left( \left\| \xi_{x_{\tau 0} v}(k\tau) - \bar{\xi}_{x_{\tau 0} \bar{v}}(k\tau) \right\|^2 \right)^{\frac{1}{2}} \right\} + \varepsilon > \varepsilon \mid x_{\tau 0} \right\} \leq \\
& \mathbb{P} \left\{ \sup_{k \in \mathbb{N}_0} \left\{ \left( \phi \left( \xi_{x_{\tau 0} v}(k\tau), \bar{\xi}_{x_{\tau 0} \bar{v}}(k\tau) \right) \right)^{\frac{1}{2}} \right\} + \varepsilon > \varepsilon \mid x_{\tau 0} \right\} \leq \\
& \quad \frac{\sqrt{\phi(x_{\tau 0}, x_{\tau 0})} + \varepsilon}{\varepsilon}.
\end{aligned}$$

In a similar way, we can prove that for any state run  $X_{x_{q 0} u}$  of  $S_q(\Sigma)$ , there exists a state run  $\xi_{x_{\tau 0} v}$  of  $S_\tau(\Sigma)$  such that the relation in (V.10) holds. ■

*Proof of Proposition 5.9:* As argued in the proof of Proposition 5.7, for any state run  $\bar{\xi}_{x_{\tau 0} v}$  of  $S_\tau(\bar{\Sigma})$  (notice that here the non-probabilistic model is excited with the same input  $v$  of  $\Sigma$ ), there exists a state run  $X_{x_{q 0} u}$  of  $S_q(\Sigma)$  such that:

$$\sup_{k \in \mathbb{N}_0} \left\| \bar{\xi}_{x_{\tau 0} v}(k\tau) - X_{x_{q 0} u}(k) \right\| \leq \varepsilon.$$

Hence, one obtains the following chain of (in)equalities:

$$\begin{aligned}
& \mathbb{P} \left\{ \sup_{0 \leq k \leq N} \|\xi_{x_{\tau_0} v}(k\tau) - X_{x_{q_0} u}(k)\| > \epsilon + \varepsilon \mid x_{\tau_0} \right\} = \\
& \mathbb{P} \left\{ \sup_{0 \leq k \leq N} \left\| \xi_{x_{\tau_0} v}(k\tau) - \bar{\xi}_{x_{\tau_0} v}(k\tau) \right. \right. \\
& \quad \left. \left. + \bar{\xi}_{x_{\tau_0} v}(k\tau) - X_{x_{q_0} u}(k) \right\| > \epsilon + \varepsilon \mid x_{\tau_0} \right\} \leq \\
& \mathbb{P} \left\{ \sup_{0 \leq k \leq N} \left\{ \left\| \xi_{x_{\tau_0} v}(k\tau) - \bar{\xi}_{x_{\tau_0} v}(k\tau) \right\| \right. \right. \\
& \quad \left. \left. + \left\| \bar{\xi}_{x_{\tau_0} v}(k\tau) - X_{x_{q_0} u}(k) \right\| \right\} > \epsilon + \varepsilon \mid x_{\tau_0} \right\} \leq \\
& \mathbb{P} \left\{ \sup_{0 \leq k \leq N} \left\{ \alpha \left( \left\| \xi_{x_{\tau_0} v}(k\tau) - \bar{\xi}_{x_{\tau_0} v}(k\tau) \right\|^q \right) \right\} > \alpha(\epsilon^q) \mid x_{\tau_0} \right\} \leq \\
& \mathbb{P} \left\{ \sup_{0 \leq k \leq N} \left\{ V \left( \xi_{x_{\tau_0} v}(k\tau), \bar{\xi}_{x_{\tau_0} v}(k\tau) \right) \right\} > \alpha(\epsilon^q) \mid x_{\tau_0} \right\}. \quad (\text{IX.23})
\end{aligned}$$

As showed for the  $\delta$ -ISS- $M_q$  Lyapunov function  $V$  in the proof of Lemma 3.8, we have:

$$\begin{aligned}
& \mathcal{L}^{v(t), v(t)} V \left( \xi_{x_{\tau_0} v}(t), \bar{\xi}_{x_{\tau_0} v}(t) \right) \leq -\kappa V \left( \xi_{x_{\tau_0} v}(t), \bar{\xi}_{x_{\tau_0} v}(t) \right) \\
& \quad + \frac{1}{2} \text{Tr} \left( \sigma \left( \xi_{x_{\tau_0} v}(t) \right)^T \partial_{x, x} V \left( \xi_{x_{\tau_0} v}(t), \bar{\xi}_{x_{\tau_0} v}(t) \right) \sigma \left( \xi_{x_{\tau_0} v}(t) \right) \right) \\
& \quad - \frac{1}{2} \text{Tr} \left( \begin{bmatrix} \sigma \left( \xi_{x_{\tau_0} v}(t) \right) \\ \sigma \left( \bar{\xi}_{x_{\tau_0} v}(t) \right) \end{bmatrix} \begin{bmatrix} \sigma^T \left( \xi_{x_{\tau_0} v}(t) \right) & \sigma^T \left( \bar{\xi}_{x_{\tau_0} v}(t) \right) \end{bmatrix} \right) \\
& \quad H(V) \left( \xi_{x_{\tau_0} v}(t), \bar{\xi}_{x_{\tau_0} v}(t) \right) \\
& \leq -\kappa V \left( \xi_{x_{\tau_0} v}(t), \bar{\xi}_{x_{\tau_0} v}(t) \right) + \frac{\alpha}{2}, \quad (\text{IX.24})
\end{aligned}$$

for any  $v \in \mathcal{U}_\tau$  and any  $x_{\tau_0} \in \mathbb{R}^n$ . Using inequalities (IX.23), (IX.24), and Theorem 1 in [20, Chapter III], one obtains the relations (V.11) and (V.12). In a similar way, we can prove that for any state run  $X_{x_{q_0} u}$  of  $S_q(\Sigma)$ , there exists a state run  $\xi_{x_{\tau_0} v}$  of  $S_\tau(\Sigma)$  such that the relations in (V.11) and (V.12) hold. ■



**Majid Zamani** (S'08–M'12) is an Assistant Professor in the Department of Electrical Engineering and Information Technology at the Technical University of Munich, Germany. He received a B.Sc. degree in Electrical Engineering in 2005 from Isfahan University of Technology, Iran, an M.Sc. degree in Electrical Engineering in 2007 from Sharif University of Technology, Iran, an MA degree in Mathematics and a Ph.D. degree in Electrical Engineering both in 2012 from University of California, Los Angeles, USA. Between September 2012 and December 2013

he was a postdoctoral researcher at the Delft Centre for Systems and Control, Delft University of Technology, Netherlands. From December 2013 to April 2014 he was an Assistant Professor in the Design Engineering Department, Delft University of Technology, Netherlands.

His research interests include verification and control of hybrid systems, embedded control software synthesis, networked control systems, and incremental properties of nonlinear control systems.



**Peyman Mohajerin Esfahani** received the B.Sc. and M.Sc. degrees in Electrical Engineering both from Sharif University of Technology, Tehran, Iran, in September 2005 and December 2007, respectively. In June 2008 he joined the Automatic Control Lab at the Swiss Federal Institute of Technology (ETH) Zurich, Switzerland, where he obtained the Ph.D. degree in January 2014.

He currently holds a dual-appointment as a postdoctoral researcher at Risk Analytics and Optimization Chair at ETH Lausanne (EPFL) and the Automatic Control Laboratory at ETH Zurich. His research interests include control synthesis, analysis, and decision making problems in uncertain, adversarial, and dynamic environments subject to information constraints, with applications to large scale systems such as power networks and industrial cyber-physical systems.



**Rupak Majumdar** received the Ph.D. degree in Computer Science from the University of California at Berkeley in 2003. He is a Scientific Director at the Max Planck Institute for Software Systems. His current research interests include verification and control of hybrid systems, automata theory and logic, software verification and program analysis. He has received the NSF CAREER award, a Sloan Foundation Fellowship, an ERC Synergy award, most influential paper awards from ACM SIGPLAN POPL and PLDI, and several best paper awards.



**Alessandro Abate** (S'02–M'08) is an Associate Professor in the Department of Computer Science at the University of Oxford (UK), and is additionally affiliated with the Delft Centre for Systems and Control at TU Delft (NL). He received a Laurea in Electrical Engineering in October 2002 from the University of Padova (IT), an MS in May 2004 and a PhD in December 2007, both in Electrical Engineering and Computer Sciences, at UC Berkeley (USA). He has been an International Fellow in the CS Lab at SRI International in Menlo Park (USA), and a

PostDoctoral Researcher at Stanford University (USA), in the Department of Aeronautics and Astronautics. From June 2009 to mid 2013 he has been an Assistant Professor at the Delft Centre for Systems and Control, TU Delft - Delft University of Technology (NL).

His research interests are in the analysis, verification, and control of probabilistic and hybrid systems, and in their general application over a number of domains, particularly in systems biology and in energy.



**John Lygeros** holds the chair of Computation and Control at the Swiss Federal Institute of Technology (ETH) Zurich, Switzerland, where he is currently serving as the Head of the Automatic Control Laboratory. He completed a B.Eng. degree in electrical engineering in 1990 and an M.Sc. degree in Systems Control in 1991, both at Imperial College of Science Technology and Medicine, London, U.K.. In 1996 he obtained a Ph.D. degree from the Electrical Engineering and Computer Sciences Department, University of California, Berkeley. During the period

1996-2000 he held a series of research appointments at the National Automated Highway Systems Consortium, Berkeley, the Laboratory for Computer Science, M.I.T., and the Electrical Engineering and Computer Sciences Department at U.C. Berkeley. Between 2000 and 2003 he was a University Lecturer at the Department of Engineering, University of Cambridge, U.K., and a Fellow of Churchill College. Between 2003 and 2006 he was an Assistant Professor at the Department of Electrical and Computer Engineering, University of Patras, Greece. In July 2006 he joined the Automatic Control Laboratory at ETH Zurich, first as an Associate Professor, and since January 2010 as Full Professor. His research interests include modeling, analysis, and control of hierarchical, hybrid, and stochastic systems, with applications to biochemical networks, automated highway systems, air traffic management, power grids and camera networks. John Lygeros is a Fellow of the IEEE, and a member of the IET and of the Technical Chamber of Greece.